

A view from infinity of the uniform infinite planar quadrangulation

N. Curien, L. Ménard, G. Miermont

January 6, 2012

Abstract

We introduce a new construction of the Uniform Infinite Planar Quadrangulation (UIPQ). Our approach is based on an extension of the Cori-Vauquelin-Schaeffer mapping in the context of infinite trees, in the spirit of previous work [11, 27, 30]. However, we release the positivity constraint on the labels of trees which was imposed in these references, so that our construction is technically much simpler. This approach allows us to prove the conjectures of Krikun [21] pertaining to the “geometry at infinity” of the UIPQ, and to derive new results about the UIPQ, among which a fine study of infinite geodesics.

1 Introduction

The purpose of this work is to develop a new approach to the Uniform Infinite Planar Quadrangulation (UIPQ), which is a model of random discrete planar geometry consisting in a cell decomposition of the plane into quadrangles, chosen “uniformly at random” among all homeomorphically distinct possibilities.

Recall that a planar map is a proper embedding of a finite connected graph in the two-dimensional sphere, viewed up to orientation-preserving homeomorphisms of the sphere. The faces are the connected components of the complement of the union of the edges. A map is a triangulation (respectively a quadrangulation) if all its faces are incident to three (respectively four) edges. A map is rooted if one has distinguished an oriented edge called the root edge. Planar maps are basic objects in combinatorics and have been extensively studied since the work of Tutte in the sixties [36]. They also appear in various areas, such as algebraic geometry [23], random matrices [37] and theoretical physics, where they have been used as a model of random geometry [2].

The latter was part of Angel and Schramm’s motivation to introduce in [4] the so-called Uniform Infinite Planar Triangulation as a model for random planar geometry. Its companion, the UIPQ, was later defined by Krikun [20] following a similar approach. One advantage of quadrangulations over triangulations is that there exists a very nice bijection between, on the one hand, rooted planar quadrangulations with n faces, and on the other hand, labeled plane trees with n edges and non-negative labels. This bijection is due to Cori and Vauquelin [13], but only reached its full extension with the work of Schaeffer [35]. See Section 2.3. This leads Chassaing and Durhuus [11] to introduce an infinite random quadrangulation of the plane, generalizing the Cori-Vauquelin-Schaeffer bijection to a construction of the random quadrangulation from an infinite random tree with non-negative labels. Ménard [30] then showed that the two constructions of [20, 11] lead to the same random object. In the Chassaing-Durhuus approach, the labels in the random infinite tree correspond to distances from the origin of the root edge in the quadrangulation, and thus information about the labels can be used to derive geometric properties such as volume growth around the root in the UIPQ [11, 27, 30].

Let us describe quickly the UIPQ with the point of view of Angel-Schramm and Krikun. If Q_n is a random rooted quadrangulation uniformly distributed over the set of all rooted quadrangulations with n faces, then we have [20]

$$Q_n \xrightarrow[n \rightarrow \infty]{(d)} Q_\infty,$$

in distribution in the sense of the *local convergence*, meaning that for every fixed $R > 0$, the combinatorial balls of Q_n with radius R and centered at the root converge in distribution as $n \rightarrow \infty$ to that of Q_∞ , see Section 2.1.2 for more details. The object Q_∞ is a random infinite rooted quadrangulation called the Uniform Infinite Planar Quadrangulation (UIPQ). The UIPQ and its sister the UIPT are fundamental objects in random geometry and have been the object of many studies. See [3, 4, 5, 20, 21, 22] and references therein.

In the present work, we give a new construction of the UIPQ from a certain random labeled tree. This is in the spirit of the “bijective” approach by Chassaing-Durhuus, but where the positivity constraint on the labels is released. Though the labels no longer correspond to distances from the root of the UIPQ, they can still be interpreted as “distances seen from the point at infinity”. In many respects, this construction is simpler than [11] because the unconditioned labeled tree has a very simple branching structure — its genealogy is that of a critical branching process conditioned on non-extinction. This simplifies certain computations on the UIPQ and enables us to derive new results easily.

Let us briefly describe our construction. We denote by T_∞ the critical geometric Galton-Watson tree conditioned to survive. This random infinite planar tree with one end has been introduced by Kesten [19] and can be built from a semi-infinite line of vertices x_0, x_1, x_2, \dots together with independent critical geometric Galton-Watson trees grafted to the left-hand side and right-hand side of each vertex x_i for $i \geq 0$, see Section 2.4. Conditionally on T_∞ , we consider a sequence of independent variables $(d_e)_{e \in E(T_\infty)}$ indexed by the edges of T_∞ which are uniformly distributed over $\{-1, 0, +1\}$. We then assign to every vertex u of T_∞ a label $\ell(u)$ corresponding to the sum of the numbers d_e along the ancestral path from u to the root x_0 of T_∞ . Given an extra Bernoulli variable $\eta \in \{0, 1\}$ independent of (T_∞, ℓ) , it is then possible to extend the classical Schaeffer construction to define a quadrangulation $\Phi((T_\infty, \ell), \eta)$ from (T_∞, ℓ) and η , see Section 2.3. The only role of η is to prescribe the orientation of the root edge in $\Phi((T_\infty, \ell), \eta)$. The random infinite rooted quadrangulation $Q_\infty = \Phi((T_\infty, \ell), \eta)$ has the distribution of the UIPQ, see Theorem 1. Moreover, the vertices of Q_∞ correspond to those of T_∞ and via this identification, Theorem 1 gives a simple interpretation of the labels: Almost surely, for any pair of vertices u, v of Q_∞ we have

$$\ell(u) - \ell(v) = \lim_{z \rightarrow \infty} \left(d_{\text{gr}}^{Q_\infty}(u, z) - d_{\text{gr}}^{Q_\infty}(v, z) \right), \quad (*)$$

where $d_{\text{gr}}^{Q_\infty}$ is the usual graph distance. The fact that the limit exists as $z \rightarrow \infty$ in $(*)$ means that the right-hand side is constant everywhere but on a finite subset of vertices of Q_∞ . Theorem 1 and its corollaries also answer positively the three conjectures raised by Krikun in [21]. Note that the existence of the limit in $(*)$ was shown in [21]. It also follows from our fine study of the geodesics and their coalescence properties in the UIPQ, see Proposition 5 and Theorem 2.

As a corollary of our new construction we study (see Theorem 3) the length of the separating cycle at a given height (seen from ∞) in the UIPQ, much in the spirit of a previous work of Krikun’s [20]. We also deduce new properties that support a conjecture of Angel & Schramm [4] (reformulated in our context) saying that the UIPQ is recurrent. Namely, we show that the distances from infinity along the random walk on the UIPQ is a recurrent process.

The paper is organized as follows. In Section 2 we introduce the construction of the UIPQ based on a random infinite labeled tree and present our main theorem. Section 3 is devoted to the proof of Theorem 1, which goes through an analysis of discrete geodesics in the UIPQ.

In particular, we establish a confluence property of geodesics towards the root (Proposition 5) and a certain uniqueness property of geodesic rays towards infinity (Theorem 2). Section 4 is devoted to the study of the scaling limits for the contour functions describing the infinite labeled tree (T_∞, ℓ) and to the proofs of two technical lemmas used to derive Theorem 2. Using our new construction we finally study separating cycles at a given height (Section 5) and random walk on the UIPQ (Section 6).

Acknowledgments: We deeply thank Jean-François Le Gall for fruitful discussions and a careful reading of a first version of this article.

2 The UIPQ and the uniform infinite labeled tree

2.1 Finite and infinite quadrangulations

Consider a proper embedding of a finite connected graph in the sphere \mathbb{S}_2 (loops and multiple edges are allowed). A *finite planar map* m is an equivalence class of such embeddings modulo orientation preserving homeomorphisms of the sphere. Let $\vec{E}(m)$ be the set of all oriented edges of m (each edge corresponds to exactly two oriented edges). A planar map is *rooted* if it has a distinguished oriented edge $e^* \in \vec{E}(m)$, which is called the root edge. If e is an oriented edge of a map we write e_- and e_+ for its origin and target vertices and \overleftarrow{e} for the reversed edge.

The set of vertices of a map m is denoted by $V(m)$. We will equip $V(m)$ with the graph distance: If v and v' are two vertices, $d_{\text{gr}}^m(v, v')$ is the minimal number of edges on a path from v to v' in m . If $v \in V(m)$, the *degree* of v is the number of oriented edges pointing towards v and is denoted by $\deg(v)$.

The *faces* of the map are the connected components of the complement of the union of its edges. The *degree* of a face is the number of edges that are incident to it, where it should be understood that an edge lying entirely in a face is incident twice to this face. A finite planar map is a *quadrangulation* if all its faces have degree 4, that is 4 incident edges. A planar map is a *quadrangulation with holes* if all its faces have degree 4, except for a number of distinguished faces which can be of arbitrary even degrees. We call these faces the *holes*, or the *boundaries* of the quadrangulation.

2.1.1 Infinite quadrangulations and their planar embeddings

Let us introduce infinite quadrangulations using the approach of Krikun [20], see also [4, 7]. For every integer $n \geq 1$, we denote by \mathbf{Q}_n the set of all rooted quadrangulations with n faces. For every pair $q, q' \in \mathbf{Q}_f = \bigcup_{n \geq 1} \mathbf{Q}_n$ we define

$$d_{\mathbf{Q}}(q, q') = \left(1 + \sup \{r : B_{\mathbf{Q}, r}(q) = B_{\mathbf{Q}, r}(q')\}\right)^{-1}$$

where, for $r \geq 1$, $B_{\mathbf{Q}, r}(q)$ is the planar map whose edges (resp. vertices) are all edges (resp. vertices) incident to a face of q having at least one vertex at distance strictly smaller than r from the root vertex e_-^* , and $\sup \emptyset = 0$ by convention. Note that $B_{\mathbf{Q}, r}(q)$ is a quadrangulation with holes.

The pair $(\mathbf{Q}_f, d_{\mathbf{Q}})$ is a metric space, we let $(\mathbf{Q}, d_{\mathbf{Q}})$ be the completion of this space. We call *infinite quadrangulations* the elements of \mathbf{Q} that are not finite quadrangulations and we denote the set of all such quadrangulations by \mathbf{Q}_∞ . Note that one can extend the function $q \in \mathbf{Q}_f \mapsto B_{\mathbf{Q}, r}(q)$ to a continuous function $B_{\mathbf{Q}, r}$ on \mathbf{Q} .

Infinite quadrangulations of the plane. An infinite quadrangulation q defines a unique infinite graph G with a root edge, together with a consistent family of planar embeddings $(B_{\mathbf{Q},r}(q), r \geq 1)$ of the combinatorial balls of G centered at the root vertex.

Conversely, any sequence q_1, q_2, \dots of rooted quadrangulations with holes, such that $q_r = B_{\mathbf{Q},r}(q_{r+1})$ for every $r \geq 1$, specifies a unique infinite quadrangulation q whose ball of radius r is q_r for every $r \geq 1$.

Definition 1. An infinite quadrangulation $q \in \mathbf{Q}_\infty$ is called a *quadrangulation of the plane* if it has one end, that is, if for any $r \geq 0$ the graph $q \setminus B_{\mathbf{Q},r}(q)$ has only one infinite connected component.

It is not hard to convince oneself that quadrangulations of the plane also coincide with equivalence classes of certain proper embeddings of infinite graphs in the plane \mathbb{R}^2 , viewed up to orientation preserving homeomorphisms. Namely these are the proper embeddings χ of locally finite planar graphs such that

- every compact subset of \mathbb{R}^2 intersects only finitely many edges of χ ,
- the connected components of the complement of the union of edges of χ in \mathbb{R}^2 are all bounded topological quadrangles.

Remark. Note that a generic element of \mathbf{Q}_∞ is not necessarily a quadrangulation of the plane. See [4, 11, 30] and the Appendix below for more details about this question.

2.1.2 The Uniform Infinite Planar Quadrangulation

Now, let Q_n be a random variable with uniform distribution on \mathbf{Q}_n . Then as $n \rightarrow \infty$, the sequence $(Q_n)_{n \geq 1}$ converges in distribution to a random variable with values in \mathbf{Q}_∞ .

Theorem ([20]). *For every $n \geq 1$, let ν_n be the uniform probability measure on \mathbf{Q}_n . The sequence $(\nu_n)_{n \geq 1}$ converges to a probability measure ν , in the sense of weak convergence in the space of probability measures on $(\mathbf{Q}, d_{\mathbf{Q}})$. Moreover, ν is supported on the set of infinite rooted quadrangulations of the plane.*

The probability measure ν is called the law of the uniform infinite planar quadrangulation (UIPQ).

2.2 Labeled trees

Throughout this work we will use the standard formalism for planar trees as found in [32]. Let

$$\mathcal{U} = \bigcup_{n=0}^{\infty} \mathbb{N}^n$$

where $\mathbb{N} = \{1, 2, \dots\}$ and $\mathbb{N}^0 = \{\emptyset\}$ by convention. An element u of \mathcal{U} is thus a finite sequence of positive integers. If $u, v \in \mathcal{U}$, uv denotes the concatenation of u and v . If v is of the form uj with $j \in \mathbb{N}$, we say that u is the *parent* of v or that v is a *child* of u . More generally, if v is of the form uw , for $u, w \in \mathcal{U}$, we say that u is an *ancestor* of v or that v is a *descendant* of u . A *rooted planar tree* τ is a (finite or infinite) subset of \mathcal{U} such that

1. $\emptyset \in \tau$ (\emptyset is called the *root* of τ),
2. if $v \in \tau$ and $v \neq \emptyset$, the parent of v belongs to τ
3. for every $u \in \mathcal{U}$ there exists $k_u(\tau) \geq 0$ such that $uj \in \tau$ if and only if $j \leq k_u(\tau)$.

A rooted planar tree can be seen as a graph, in which an edge links two vertices u, v such that u is the parent of v or vice-versa. This graph is of course a tree in the graph-theoretic sense, and has a natural embedding in the plane, in which the edges from a vertex u to its children $u_1, \dots, u_{k_u}(\tau)$ are drawn from left to right.

We let $|u|$ be the length of the word u . The integer $|\tau|$ denotes the number of edges of τ and is called the size of τ . A *spine* in a tree τ is an infinite sequence u_0, u_1, u_2, \dots in τ such that $u_0 = \emptyset$ and u_i is the parent of u_{i+1} for every $i \geq 0$. If a and b are two vertices of a tree τ , we denote the set of vertices along the unique geodesic path going from a to b in τ by $[[a, b]]$.

A *rooted labeled tree* (or spatial tree) is a pair $\theta = (\tau, (\ell(u))_{u \in \tau})$ that consists of a rooted planar tree τ and a collection of integer labels assigned to the vertices of τ , such that if $u, v \in \tau$ and v is a child of u , then $|\ell(u) - \ell(v)| \leq 1$. For every $l \in \mathbb{Z}$, we denote by $\mathbf{T}^{(l)}$ the set of labeled trees for which $\ell(\emptyset) = l$, and $\mathbf{T}_\infty^{(l)}$, resp. $\mathbf{T}_f^{(l)}$, resp. $\mathbf{T}_n^{(l)}$, are the subsets of $\mathbf{T}^{(l)}$ consisting of the infinite trees, resp. finite trees, resp. trees with n edges. If $\theta = (\tau, \ell)$ is a labeled tree, $|\theta| = |\tau|$ is the size of θ .

If θ is a labeled tree and $h \geq 0$ is an integer, we denote the labeled subtree of θ consisting of all vertices of θ and their labels up to height h by $B_{\mathbf{T}, h}(\theta)$. For every pair θ, θ' of labeled trees define

$$d_{\mathbf{T}}(\theta, \theta') = (1 + \sup \{h : B_{\mathbf{T}, h}(\theta) = B_{\mathbf{T}, h}(\theta')\})^{-1}.$$

One easily checks that $d_{\mathbf{T}}$ is a distance on the set of all labeled trees, which turns this set into a separable and complete metric space.

In the rest of this work we will mostly be interested in the following set of infinite trees. We let \mathcal{S} be the set of all labeled trees (τ, ℓ) in $\mathbf{T}_\infty^{(0)}$ such that

- τ has exactly one spine, which we denote by $\emptyset = S_\tau(0), S_\tau(1), S_\tau(2), \dots$
- $\inf_{i \geq 0} \ell(S_\tau(i)) = -\infty$.

If $\theta = (\tau, \ell) \in \mathcal{S}$, the spine then splits τ in two parts, which we call the left and right parts, and every vertex $S_\tau(n)$ of the spine determines a subtree of τ to its left and to its right. These are denoted by $L_n(\theta), R_n(\theta)$, formally,

$$\begin{aligned} L_n(\theta) &= \{v \in \mathcal{U} : S_\tau(n)v \in \tau, S_\tau(n)v \prec S_\tau(n+1)\} \\ R_n(\theta) &= \{v \in \mathcal{U} : S_\tau(n)v \in \tau, S_\tau(n+1) \prec S_\tau(n)v\} \cup \{\emptyset\}, \end{aligned}$$

where $u \prec v$ denotes the lexicographical order on \mathcal{U} . The subtrees $L_n(\theta), R_n(\theta)$ naturally inherit the labels from θ , so that we really see $L_n(\theta), R_n(\theta)$ as elements of $\mathbf{T}_f^{(X_n(\theta))}$, where $X_n(\theta) = \ell(S_\tau(n))$ is the label of the n -th vertex of the spine. We can of course reconstruct the tree θ from the sequence $(L_n(\theta), R_n(\theta))_{n \geq 0}$. In the sequel, we will often write $S(n), X_n, L_n, R_n$ instead of $S_\tau(n), X_n(\theta), L_n(\theta), R_n(\theta)$ when there is no ambiguity on the underlying labeled tree.

2.3 The Schaeffer correspondence

One of the main tools for studying random quadrangulations is a bijection initially due to Cori & Vauquelin [13], and that was much developed by Schaeffer [35]. It establishes a one-to-one correspondence between rooted and pointed quadrangulations with n faces, and pairs consisting of a labeled tree of $\mathbf{T}_n^{(0)}$ and an element of $\{0, 1\}$. Let us describe this correspondence and its extension to infinite quadrangulations.

2.3.1 From trees to quadrangulations

A rooted and *pointed* quadrangulation is a pair $\mathbf{q} = (q, \rho)$ where q is a rooted quadrangulation and ρ is a distinguished vertex of q . We write \mathbf{Q}_n^\bullet for the set of all rooted and pointed quadrangulations with n faces. We first describe the mapping from labeled trees to quadrangulations.

Let $\theta = (\tau, \ell)$ be an element of $\mathbf{T}_n^{(0)}$. We view τ as embedded in the plane. A *corner* of a vertex in τ is an angular sector formed by two consecutive edges in clockwise order around this vertex. Note that a vertex of degree k in τ has exactly k corners. If c is a corner of τ , $\mathcal{V}(c)$ denotes the vertex incident to c . By extension, the label $\ell(c)$ of a corner c is the label of $\mathcal{V}(c)$.

The corners are ordered clockwise cyclically around the tree in the so-called *contour order*. If we view τ as a planar map with one face, then τ can be seen as a polygon with $2n$ edges that are glued by pairs, and the contour order is just the usual cyclic order of the corners of this polygon. We fix the labeling by letting $(c_0, c_1, c_2, \dots, c_{2n-1})$ be the sequence of corners visited during the contour process of τ , starting from the corner c_0 incident to \emptyset that is located to the left of the oriented edge going from \emptyset to 1 in τ . We extend this sequence of corners into a sequence $(c_i, i \geq 0)$ by periodicity, letting $c_{i+2n} = c_i$. For $i \in \mathbb{Z}_+$, the *successor* $\mathcal{S}(c_i)$ of c_i is the first corner c_j in the list $c_{i+1}, c_{i+2}, c_{i+3}, \dots$ of label $\ell(c_j) = \ell(c_i) - 1$, if such a corner exists. In the opposite case, the successor of c_i is an extra element ∂ , not in $\{c_i, i \geq 0\}$.

Finally, we construct a new graph as follows. Add an extra vertex ρ in the plane, that does not belong to (the embedding of) τ . For every corner c , draw an arc between c and its successor if this successor is not ∂ , or draw an arc between c and ρ if the successor of c is ∂ . The construction can be made in such a way that the arcs do not cross. After the interior of the edges of τ has been removed, the resulting embedded graph, with vertex set $\tau \cup \{\rho\}$ and edges given by the newly drawn arcs, is a quadrangulation q . In order to root this quadrangulation, we consider some extra parameter $\eta \in \{0, 1\}$. If $\eta = 0$, the root of q is the arc from c_0 to its successor, oriented in this direction. If $\eta = 1$ then the root of q is the same edge, but with opposite orientation. We let $q = \Phi(\theta, \eta) \in \mathbf{Q}_n^\bullet$ (q comes naturally with the distinguished vertex ρ).

Theorem (Theorem 4 in [12]). *The mapping $\Phi : \mathbf{T}_n^{(0)} \times \{0, 1\} \longrightarrow \mathbf{Q}_n^\bullet$ is a bijection. If $q = \Phi((\tau, \ell), \eta)$ then for every vertex v of q not equal to ρ , one has*

$$d_{\text{gr}}^q(v, \rho) = \ell(v) - \min_{u \in \tau} \ell(u) + 1, \quad (1)$$

where we recall that every vertex of q not equal to ρ is identified to a vertex of τ .

Note that (1) can also be rewritten as

$$\ell(v) = d_{\text{gr}}^q(v, \rho) - d_{\text{gr}}^q(e_\pm^*, \rho), \quad v \in V(q), \quad (2)$$

where


$$e_\pm^* = \emptyset = \begin{cases} e_-^* & \text{if } d_{\text{gr}}^q(e_-^*, \rho) - d_{\text{gr}}^q(e_+^*, \rho) = -1 \\ e_+^* & \text{if } d_{\text{gr}}^q(e_-^*, \rho) - d_{\text{gr}}^q(e_+^*, \rho) = 1 \end{cases}$$

Hence, these labels can be recovered from the pointed quadrangulation (q, ρ) . This is of course not surprising since the function $\Phi : \mathbf{T}_n^{(0)} \times \{0, 1\} \rightarrow \mathbf{Q}_n^\bullet$ is invertible (see the next section for the description of the inverse mapping).

Infinite case. We now aim at extending the construction of Φ to elements of \mathcal{S} . Let $(\tau, (\ell(u))_{u \in \tau}) \in \mathcal{S}$. Again, we consider an embedding of τ in the plane, with isolated vertices. This is always possible (since τ is locally finite). The notion of a corner is unchanged in this setting, and there is still a notion of clockwise contour order for the corners of τ , this order being now a total order, isomorphic to (\mathbb{Z}, \leq) , rather than a cyclic order. We consider the sequence $(c_0^{(L)}, c_1^{(L)}, c_2^{(L)}, \dots)$ of corners visited by the contour process of the left side of the tree

$$c_i = \begin{cases} c_i^{(L)} & \text{if } i \geq 0 \\ c_{-i}^{(R)} & \text{if } i < 0. \end{cases}$$

In a similar way as before, the embedded graph q is rooted at the edge emerging from the distinguished corner c_0 of \emptyset , that is, the edge between c_0 and its successor $\mathcal{S}(c_0)$. The direction of the edge is given by an extra parameter $\eta \in \{0, 1\}$, similarly as above.



Proposition 2. *The resulting embedded graph q is an infinite quadrangulation of the plane, and the extended mapping $\Phi : \mathcal{S} \cup \mathbf{T}_f^{(0)} \rightarrow \mathbf{Q}$ is continuous.*

7

there exists a corner c_j with $j < i$ such that the vertex incident to c_j belongs to the spine $\{S_\tau(0), S_\tau(1), \dots\}$, and $\min_{j \leq k \leq i} \ell(c_k) < \ell(c_i) - 1$. Therefore, for every $k \leq j$, the successor of c_k is not c_i .

Together with the fact that every vertex has a number of successors equal to its degree, this shows that the embedded graph q is locally finite, in the sense that every vertex is incident to a finite number of edges. The fact that every face of q is a quadrangle is then a consequence of the construction of the arcs, as proved e.g. in [12]. It remains to show that q can be properly embedded in the plane, that is, has one end. This comes from the construction of the edges and the fact that τ has only one end. The details are left to the reader.

To prove the continuity of Φ , let $\theta_n = (\tau_n, \ell_n)$ be a sequence in $\mathcal{S} \cup \mathbf{T}_f^{(0)}$ converging to $\theta = (\tau, \ell) \in \mathcal{S} \cup \mathbf{T}_f^{(0)}$. If $\theta \in \mathbf{T}_f^{(0)}$ then $\theta_n = \theta$ for every n large enough, so the fact that $\Phi(\theta_n) \rightarrow \Phi(\theta)$ is obvious. So let us assume that $\theta \in \mathcal{S}$, with spine vertices $S_\tau(0), S_\tau(1), \dots$. Let $R > 0$ be an integer, and let $l(R)$ be the minimal label of a vertex in $B_{\mathbf{T}, R}(\theta)$. Since $\inf(\ell(S_\tau(i))) = -\infty$, we can define $f(R) > R$ as the first $i \geq 1$ such that $\ell(S_\tau(i)) = l(R) - 2$. If c is a corner in the subtree of τ above $S_\tau(f(R))$, then the successor of c cannot be in $B_{\mathbf{T}, R}(\theta)$. Indeed, if $\ell(c) \geq l(R) - 1$ then the successor of c has to be also in the subtree of τ above $S_\tau(f(R))$, while if $\ell(c) < l(R) - 1$, then this successor also has label $< l(R) - 1$, and thus cannot be in $B_{\mathbf{T}, R}(\theta)$ by definition. Similarly, c cannot be the successor of any corner in $B_{\mathbf{T}, R}(\theta)$, as these successors necessarily are in the subtree of τ below $S_\tau(f(R))$.

Now, for every n large enough, it holds that $B_{\mathbf{T}, f(R)}(\theta_n) = B_{\mathbf{T}, f(R)}(\theta)$, from which we obtain that the maps formed by the arcs incident to the vertices of $B_{\mathbf{T}, R}(\theta) = B_{\mathbf{T}, R}(\theta_n)$ are the same, and moreover, no extra arc constructed in θ_n or θ is incident to a vertex of $B_{\mathbf{T}, R}(\theta) = B_{\mathbf{T}, R}(\theta_n)$. Letting $r > 0$ and choosing R so that all the edges of $B_{\mathbf{Q}, r}(\Phi(\theta))$ appear as arcs incident to vertices of $B_{\mathbf{T}, R}(\theta)$, we obtain that $B_{\mathbf{Q}, r}(\Phi(\theta)) = B_{\mathbf{Q}, r}(\Phi(\theta_n))$ for n large enough. Therefore, we get that $\Phi(\theta_n) \rightarrow \Phi(\theta)$, as desired. \square

The vertex set of q is precisely τ , so that the labels ℓ on τ induce a labeling of the vertices of q . In the finite case, we saw earlier in (2) that these labels could be recovered from the pointed quadrangulation obtained from a finite labeled tree. In our infinite setting, this is much less obvious: Intuitively the distinguished vertex ρ of the finite case is “lost at infinity”.

Bounds on distances. We will see later that when the infinite labeled tree has a special distribution corresponding *via* the Schaeffer correspondence Φ to the UIPQ, then the labels have a natural interpretation in terms of distances in the infinite quadrangulation. In general if an infinite quadrangulation q is constructed from a labeled tree $\theta = (\tau, \ell)$ in \mathcal{S} , every pair $\{u, v\}$ of neighboring vertices in q satisfies $|\ell(u) - \ell(v)| = 1$ and thus for every $a, b \in q$ linked by a geodesic $a = a_0, a_1, \dots, a_{d_{\text{gr}}^q(a, b)} = b$ we have the crude bound

$$d_{\text{gr}}^q(a, b) = \sum_{i=1}^{d_{\text{gr}}^q(a, b)} |\ell(a_i) - \ell(a_{i-1})| \geq \left| \sum_{i=1}^{d_{\text{gr}}^q(a, b)} \ell(a_i) - \ell(a_{i-1}) \right| = |\ell(a) - \ell(b)|. \quad (3)$$

A better upper bound is given by the so-called *cactus bound*

$$d_{\text{gr}}^q(a, b) \geq \ell(a) + \ell(b) - 2 \min_{v \in \llbracket a, b \rrbracket} \ell(v), \quad (4)$$

where we recall that $\llbracket a, b \rrbracket$ represents the geodesic line in τ between a and b . This bound is proved in [14] in the context of finite trees and quadrangulations, but remains valid here without change. The idea goes as follows: let w be of minimal label on $\llbracket a, b \rrbracket$, and assume $w \notin \{a, b\}$ to avoid trivialities. Removing w breaks the tree τ into two connected parts, containing respectively a and b . Now a path from a to b has to “pass over” w using an arc between a corner (in the first component) to its successor (in the other component), and this can only happen by visiting a vertex with label less than $\ell(w)$. Using (3) we deduce that this path has length at least $\ell(a) - \ell(w) + \ell(b) - \ell(w)$, as wanted.

2.3.2 From quadrangulations to trees

We saw that the Schaeffer mapping $\mathbf{T}_n^{(0)} \times \{0,1\} \longrightarrow \mathbf{Q}_n^\bullet$ is in fact a bijection. We now describe the reverse construction. The details can be found in [12]. Let (q, ρ) be a finite rooted quadrangulation given with a distinguished vertex $\rho \in V(q)$. We define a labeling ℓ of the vertices of the quadrangulation by setting

$$\ell(v) = d_{\text{gr}}^q(v, \rho), \quad v \in V(q).$$

Since the map q is bipartite, if u, v are neighbors in q then $|\ell(u) - \ell(v)| = 1$. Thus the faces of q can be decomposed into two subsets: The faces such that the labels of the vertices listed in clockwise order are $(i, i+1, i, i+1)$ for some $i \geq 0$ or those for which these labels are $(i, i+1, i+2, i+1)$ for some $i \geq 0$. We then draw on top of the quadrangulation an edge in each face according to the rules given by the figure below.




Figure 2: Rules for the reverse Schaeffer construction.

The graph τ formed by the edges added in the faces of q is a spanning tree of $q \setminus \{\rho\}$, see [12, Proposition 1]. This tree comes with a natural embedding in the plane, and we root τ according to the following rules (see Fig.3):

- If $\ell(e_-^*) > \ell(e_+^*)$ then we root τ at the corner incident to the edge e^* on e_-^* ,
- otherwise we root τ at the corner incident to the edge e^* on e_+^* ,

Finally, we shift the labeling of τ inherited from the labeling on $V(q) \setminus \{\rho\}$ by the label of the root of τ ,

$$\tilde{\ell}(u) = \ell(u) - \ell(\text{root}), \quad u \in \tau.$$

Then we have [12, Proposition 1]

$$\Phi^{-1}((q, \rho)) = ((\tau, \tilde{\ell}), \mathbf{1}_{\ell(e_+^*) > \ell(e_-^*)}).$$

Infinite case. If q is a (possibly infinite) quadrangulation and $\ell : V(q) \longrightarrow \mathbb{Z}$, is a labeling of the vertices of q such that for any neighboring vertices u, v we have $|\ell(u) - \ell(v)| = 1$, then a graph can be associated to (q, ℓ) by the device we described above. This graph could contain cycles and is not a tree in the general case.

However, suppose that the infinite quadrangulation q is constructed as the image under Φ of a labeled tree $\theta = (\tau, \ell) \in \mathcal{S}$ and an element of $\{0,1\}$. Then, with the usual identification of $V(q)$ with τ , the labeling of $V(q)$ inherited from the labeling ℓ of τ satisfies $|\ell(u) - \ell(v)| = 1$ for any $u, v \in V(q)$. An easy adaptation of the argument of [11, Property 6.2] then shows that the faces of q are in one-to-one correspondence with the edges of τ and that the edges constructed on top of each face of q following the rules of Fig. 2 exactly correspond to the edges of τ . In other words, provided that q is constructed from $\theta = (\tau, \ell)$ then the graph constructed on top of q using the labeling ℓ is exactly τ . The rooting of τ is also recovered from q and ℓ by the same procedure as in the finite case.




Figure 3: Illustration of the rooting of the plane tree τ

2.4 The uniform infinite labeled tree

For every integer $l > 0$, we denote by ρ_l the law of the Galton-Watson tree with geometric offspring distribution with parameter $1/2$, labeled according to the following rules. The root has label l and every other vertex has a label chosen uniformly in $\{m-1, m, m+1\}$ where m is the label of its parent, these choices being made independently for every vertex. Otherwise said, for every tree $\theta \in \mathbf{T}^{(l)}$, $\rho_l(\theta) = \frac{1}{2}12^{-|\theta|}$.

Definition 3. Let $\theta = (T_\infty, (\ell(u))_{u \in T_\infty})$ be a random variable with values in $(\mathbf{T}^{(0)}, d_{\mathbf{T}})$ whose distribution μ is described by the following properties

1. θ belongs to \mathcal{S} almost surely,
2. the process $(\ell(S_{T_\infty}(n)))_{n \geq 0}$ is a random walk with independent uniform steps in $\{-1, 0, 1\}$,
3. conditionally given $(\ell(S_{T_\infty}(n)))_{n \geq 0} = (x_n)_{n \geq 0}$, the sequence $(L_n(\theta))_{n \geq 0}$ of subtrees of θ attached to the left side of the spine and the sequence $(R_n(\theta))_{n \geq 0}$ of subtrees attached to the right side of the spine form two independent sequences of independent labeled trees distributed according to the measures $(\rho_{x_n})_{n \geq 0}$.

In other words, if $\theta = (T_\infty, \ell)$ is distributed according to μ then the structure of the tree T_∞ is given by an infinite spine and independent critical geometric Galton-Watson trees grafted on the left and right of each vertex of the spine. Conditionally on T_∞ the labeling is given by independent variables uniform over $\{-1, 0, +1\}$ assigned to each edge of T_∞ , which represent the label increments along the different edges, together with the boundary condition $\ell(\emptyset) = 0$.

The random infinite tree T_∞ , called the critical geometric Galton-Watson tree *conditioned to survive*, was constructed in [19, Lemma 1.14] as the limit of critical geometric Galton-Watson conditioned to survive up to level n , as $n \rightarrow \infty$. To make the link between the classical construction of T_∞ (see e.g. [29, Chapter 12]) and the one provided by the last definition, note the following equality in distribution

$$1 + G + G' \stackrel{(d)}{=} \hat{G},$$

where G, G', \hat{G} are independent random variables such that G, G' are geometric of parameter $1/2$ and \hat{G} is a size-biased geometric $1/2$ variable, that is $\mathbb{P}(\hat{G} = k) = k\mathbb{P}(G = k) = k2^{-(k+1)}$.

The law μ can also be seen as the law of a uniform infinite element of \mathcal{S} , as formalized by the following statement.

Theorem ([19]). *For every $n \geq 1$, let μ_n be the uniform probability measure on $\mathbf{T}_n^{(0)}$. Then the sequence $(\mu_n)_{n \in \mathbb{N}}$ converges weakly to μ in the space of Borel probability measures on $(\mathbf{T}^{(0)}, d_{\mathbf{T}})$.*

Proof. It is a standard result [25] that the distribution of a uniformly chosen planar tree T_n with n edges is the same as the distribution of a critical Galton-Watson tree with geometric offspring distribution conditioned on the total progeny to be $n+1$. The convergence in distribution of T_n towards T_∞ in the sense of $d_{\mathbf{T}}$ then follows from [19, Lemma 1.14], see also [29]. An analogous result holds for the uniform labeled trees since the labeling is given by independent variables uniform over $\{-1, 0, +1\}$ assigned to each edge of the trees. \square

2.5 The main result

We are now ready to state our main result. Recall that ν is the law of the UIPQ as defined in Theorem 2.1.2. Let also $\mathcal{B}(1/2)$ be the Bernoulli law $(\delta_0 + \delta_1)/2$, and recall the Schaeffer correspondence $\Phi : \mathcal{S} \times \{0, 1\} \rightarrow \mathbf{Q}$. In the following statement, if q is an element of \mathbf{Q}_∞ , and $f : V(q) \rightarrow \mathbb{Z}$ is a function on $V(q)$, we say that $\lim_{z \rightarrow \infty} f(z) = l$ if f is equal to l everywhere but on a finite subset of $V(q)$.

Theorem 1. *The probability measure ν is the image of $\mu \otimes \mathcal{B}(1/2)$ under the mapping Φ :*

$$\nu = \Phi_*(\mu \otimes \mathcal{B}(1/2)) \quad (5)$$

Moreover, if $(\theta = (T_\infty, \ell), \eta)$ has distribution $\mu \otimes \mathcal{B}(1/2)$ and $Q_\infty = \Phi(\theta, \eta)$, then, with the usual identification of the vertices of Q_∞ with the vertices of θ , one has, almost surely,

$$\ell(u) - \ell(v) = \lim_{z \rightarrow \infty} \left(d_{\text{gr}}^{Q_\infty}(u, z) - d_{\text{gr}}^{Q_\infty}(v, z) \right), \quad \forall u, v \in V(Q_\infty). \quad (6)$$

Let us make some comments about this result. The first part of the statement is easy: Since Φ is continuous from $(\mathcal{S} \cup \mathbf{T}_f^{(0)}) \times \{0, 1\}$ to \mathbf{Q} and since, if ν_n is the uniform law on \mathbf{Q}_n , one has

$$\nu_n = \Phi(\mu_n \otimes \mathcal{B}(1/2)),$$

and one obtains (5) simply by passing to the limit $n \rightarrow \infty$ in this identity using Theorems 2.1.2 and 2.4. To be completely accurate, the mapping Φ in the previous display should be understood as taking values in \mathbf{Q}_n rather than \mathbf{Q}_n^\bullet , simply by “forgetting” the distinguished vertex arising in the construction of Schaeffer’s bijection.

The rest of the statement is more subtle, and says that the labels, inherited on the vertices of Q_∞ in its construction from a labeled tree (T_∞, ℓ) distributed according to μ , can be recovered as a measurable function of Q_∞ . This is not obvious at first, because a formula such as (2) is lacking in the infinite setting. It should be replaced by the asymptotic formula (6), which specializes to

$$\ell(u) = \lim_{z \rightarrow \infty} \left(d_{\text{gr}}^{Q_\infty}(z, u) - d_{\text{gr}}^{Q_\infty}(z, e_\pm^*) \right), \quad u \in V(Q_\infty), \quad (7)$$

where

$$e_\pm^* = \begin{cases} e_-^* & \text{if } \lim_{z \rightarrow \infty} (d_{\text{gr}}^{Q_\infty}(e_-^*, z) - d_{\text{gr}}^{Q_\infty}(e_+^*, z)) = -1 \\ e_+^* & \text{if } \lim_{z \rightarrow \infty} (d_{\text{gr}}^{Q_\infty}(e_-^*, z) - d_{\text{gr}}^{Q_\infty}(e_+^*, z)) = 1 \end{cases}. \quad (8)$$

Of course, the fact that the limits in (6) and (8) exist is not obvious and is part of the statement. This was first observed by Krikun in [21], and will be derived here by different methods. Note that the vertex e_\pm^* corresponds to the root vertex \emptyset of T_∞ in the natural identification of vertices of Q_∞ with vertices of T_∞ .

In particular, the fact that the labels are measurable with respect to Q_∞ entails that (θ, η) can be recovered as a measurable function of Q_∞ . Indeed, by the discussion at the end of Section 2.3.2, the tree T_∞ can be reconstructed from Q_∞ and the labeling ℓ . The Bernoulli variable η is also recovered by (8). This settles the three conjectures proposed by Krikun in [21].

The proof of (6) depends on certain properties of geodesics in the UIPQ that we derive in the next section. Before this, we give another view on our result in terms of asymptotic geometry of the UIPQ.

2.5.1 Gromov compactification of the UIPQ

Let (X, d) be a locally compact metric space. The set $C(X)$ of real-valued continuous functions on X is endowed with the topology of uniform convergence on every compact set of X . One defines an equivalence relation on $C(X)$ by declaring two functions equal if they differ by an additive constant and the associated quotient space endowed with the quotient topology is denoted by $C(X)/\mathbb{R}$. Following [18], one can embed the original space X in $C(X)/\mathbb{R}$ using the injection

$$i: \begin{array}{ccccc} X & \longrightarrow & C(X) & \longrightarrow & C(X)/\mathbb{R} \\ x & \longmapsto & d_x = d(x, \cdot) & \longmapsto & \overline{d_x}. \end{array}$$

The *Gromov compactification* of X is then the closure of $i(X)$ in $C(X)/\mathbb{R}$. The Gromov boundary ∂X of X is composed of the points in the closure of $i(X)$ in $C(X)/\mathbb{R}$ which are not already in $i(X)$. The points in ∂X are called *horofunctions*, see [18].

Applying this discussion to the case where $(X, d) = (V(Q_\infty), d_{\text{gr}}^{Q_\infty})$, we can immediately rephrase the last part of Theorem 1 as follows.

Corollary 4. *Almost surely, the Gromov boundary ∂Q_∞ of the UIPQ consists of only one point which is $\bar{\ell}$, the equivalence class of ℓ up to additive constants.*

3 Geodesics in the UIPQ

Geodesics. If $G = (V, E)$ is a graph, a *chain* or *path* in G is a (finite or infinite) sequence of vertices $\gamma = (\gamma(0), \gamma(1), \dots)$ such that for every $i \geq 0$, the vertices $\gamma(i)$ and $\gamma(i+1)$ are linked by an edge of the graph. Such a chain is called a *geodesic* if for every $i, j \geq 0$, the graph distance d_{gr}^G between $\gamma(i)$ and $\gamma(j)$ is equal to $|j - i|$. A *geodesic ray* emanating from x is an infinite geodesic starting at $x \in V$.

We will establish two properties of the geodesics in the UIPQ: A confluence property towards the root (Section 3.1) and a confluence property towards infinity (Section 3.2). These two properties are reminiscent of the work of Le Gall on geodesics in the Brownian Map [26]. Put together they yield the last part (6) of Theorem 1.

3.1 Confluent geodesics to the root

Let Q_∞ be distributed according to ν (see Theorem 2.1.2) and x be a vertex in Q_∞ . For every $R \geq 0$, we want to show that (with probability 1) it is possible to find $R' \geq R$ and a family of geodesics $\gamma_R^z, z \notin B_{\mathbf{Q}, R'}(Q_\infty)$ linking x to z respectively, such that for every $z, z' \notin B_{\mathbf{Q}, R'}(Q_\infty)$,

$$\gamma_R^z(i) = \gamma_R^{z'}(i), \quad \text{for every } i \in \{0, 1, \dots, R\}.$$

In other words, all of these geodesics start with a common initial segment, independently of the target vertex z .

To this end, we need the construction by Chassaing-Durhuus [11] of the UIPQ, which we briefly recall. Let $l \geq 1$ and set $\overline{\mathbf{T}}^{(l)}$ be the subset of $\mathbf{T}^{(l)}$ consisting of all trees $\theta = (\tau, \ell)$ such that $\ell(v) \geq 1$ for every $v \in \tau$. Elements of $\overline{\mathbf{T}}^{(l)}$ are called l -well-labeled trees, and just well-labeled trees if $l = 1$. We let $\overline{\mathbf{T}}_n^{(l)}$ (resp. $\overline{\mathbf{T}}_\infty^{(l)}$) be the set of all l -well-labeled trees with n edges (resp. of infinite l -well-labeled trees).

Let $\overline{\mu}_n$ be the uniform distribution on $\overline{\mathbf{T}}_n^{(1)}$. Let also $\overline{\mathcal{S}}$ be the set of all trees $\theta = (\tau, \ell) \in \overline{\mathbf{T}}_\infty^{(1)}$ such that

- the tree τ has a unique spine, and
- for every $R \in \mathbb{N}$, the set $\{v \in \tau : \ell(v) \leq R\}$ is finite.

Proposition 5 ([11]). *The sequence $(\overline{\mu}_n)_{n \geq 1}$ converges weakly to a limiting probability law $\overline{\mu}$, in the space of Borel probability measures on $(\overline{\mathbf{T}}^{(1)}, d_{\mathbf{T}})$. Moreover, we have $\overline{\mu}(\overline{\mathcal{S}}) = 1$.*

The exact description of $\overline{\mu}$ is not important for our concerns, and can be found in [11]. The Schaeffer correspondence $\overline{\Phi}$ can be defined on $\overline{\mathcal{S}}$. Let us describe quickly this correspondence. Details can be found in [11], see also [27, 30].

Let $\theta = (\tau, \ell)$ be an element of $\overline{\mathcal{S}}$. We start by embedding τ in the plane in such a way that there are no accumulation points (which is possible since τ is locally finite). We add an extra vertex ∂ in the plane, not belonging to the embedding of τ . Then, we let $(c_i^{(L)}, i \geq 0)$ and $(c_i^{(R)}, i \geq 0)$ be the sequence of corners visited in contour order on the left and right sides, starting with the root corner of τ . We let, for $i \in \mathbb{Z}$,

$$c_i = \begin{cases} c_i^{(L)} & \text{if } i \geq 0 \\ c_{-i}^{(R)} & \text{if } i < 0 \end{cases}.$$

We now define the notion of successor. If the label of $\mathcal{V}(c_i)$ is 1, then the successor of the corner c_i is ∂ . Otherwise, the successor of c_i is the first corner c_j in the infinite list $\{c_{i+1}, c_{i+2}, \dots\} \cup \{\dots, c_{i-2}, c_{i-1}\}$ such that $\ell(c_j) = \ell(c_i) - 1$. The successor of any corner c_i with $\ell(c_i) \geq 2$ exists because of the labeling constraints, and the definition of $\overline{\mathcal{S}}$.

The end of the construction is similar to Section 2.3: We draw an edge between each corner and its successor and then remove all the edges of the embedding of τ . The new edges can be drawn in such a way that the resulting embedded graph is proper and represents an infinite quadrangulation of the plane. We denote this quadrangulation by $\overline{\Phi}(\theta)$ and root it at the arc from ∂ to c_0 . Note that in this construction, we do not need to introduce an extra parameter η to determine the orientation of the root. Moreover the non-negative labels ℓ have the following interpretation in terms of distances in $\overline{\Phi}(\theta)$. For every $u \in \tau$,

$$\ell(u) = d_{\text{gr}}^{\overline{\Phi}(\theta)}(\partial, u), \quad (9)$$

with the identification of the vertices of $\overline{\Phi}(\theta)$ with $\tau \cup \{\partial\}$.

Proposition 6 ([11],[30]). *It holds that*

$$\nu = \overline{\Phi}_* \overline{\mu},$$

that is, the UIPQ follows the distribution of $\overline{\Phi}(\theta)$, where θ is random with distribution $\overline{\mu}$.

Notice that the mapping $\overline{\Phi} : \overline{\mathcal{S}} \rightarrow \mathbf{Q}$ is injective. Its inverse function $\overline{\Phi}^{-1} : \overline{\Phi}(\overline{\mathcal{S}}) \rightarrow \overline{\mathbf{T}}^{(1)}$ is described in a similar manner as in Section 2.3.2: Given the quadrangulation $q = \overline{\Phi}(\tau, \ell)$, we recover the labeling ℓ over $V(q) \setminus \{\partial\}$ by (9) and $\ell(\partial) = 0$. Note that ∂ is always the origin of the root edge of q . We then apply the same device as for Φ^{-1} , that is, separating the faces of q into two kinds and adding an edge on top of them according to Fig. 2. The resulting graph is τ and is rooted at the corner incident to the root edge of q . One can check that the mapping $\overline{\Phi}^{-1}$

is continuous, i.e. that for every $h > 0$, the neighborhood $B_{\mathbf{T},h}(\tau, \ell)$ is determined by $B_{\mathbf{Q},r}(q)$ as soon as r is large enough. Thus if Q_∞ is distributed according to μ , one can define a labeled tree (τ, ℓ) distributed according to $\bar{\mu}$ as a measurable function of Q_∞ such that $Q_\infty = \bar{\Phi}(\tau, \ell)$.

From this construction, it is possible to specify a particular infinite geodesic (or *geodesic ray*) starting from e_-^* . Namely, if $(c_i)_{i \in \mathbb{Z}}$ is the contour sequence of τ , for every $i \geq 1$, let

$$d(i) = \min\{j \leq 0 : \ell(c_j) = i\},$$

which is finite by definition of $\bar{\mathcal{S}}$. Then there is an arc between $c_{d(i+1)}$ and $c_{d(i)}$ for every $i \geq 1$, as well as an arc from $c_{d(1)}$ to ∂ , and the path $(\partial, \mathcal{V}(c_{d(1)}), \mathcal{V}(c_{d(2)}), \dots)$ is a geodesic ray. We call it the *distinguished geodesic ray* of Q_∞ , and denote it by Γ , see Fig. 4.

Lemma 7. *For every $R \geq 0$, there exists $R' \geq R$ such that every $z \in V(Q_\infty) \setminus B_{\mathbf{Q},R'}(Q_\infty)$ can be joined to ∂ by a geodesic chain γ such that $\gamma(i) = \Gamma(i)$ for every $i \in \{0, 1, 2, \dots, R\}$.*

Proof. Let Q_∞ be distributed according to ν and set $(\tau, \ell) = \bar{\Phi}^{-1}(Q_\infty)$. Finally define Γ as above. Define

$$R' = \max_{d(R) \leq i \leq g(R)} \ell(c_i),$$

where $d(R)$ is defined above, and

$$g(i) = \max\{j \geq 0 : \ell(c_j) = i\}.$$

Let z be a vertex of Q_∞ , not in $B_{\mathbf{Q},R'}(Q_\infty)$, and let c_j be any corner incident to z . Then j cannot be in $[d(R), g(R)]$ since by definition $\ell(c_j) = d_{\text{gr}}^{Q_\infty}(\partial, z) > R' \geq \ell(c_i)$ for any $i \in [d(R), g(R)]$. Now, let γ be the geodesic defined as the path starting at c_j , and following the arcs from c_j to its successor corner, then from this corner to its successor, and so on until it reaches ∂ . These geodesics have the desired property, see Fig. 4. Note that if $j > 0$, that is, if c_j lies on the left side of τ , then necessarily all vertices in the geodesic γ with label less than or equal to R have to lie on the right-hand side of τ . See Fig. 4. \square

3.2 Coalescence of proper geodesics rays to infinity

With the notation of Theorem 1, let $(\theta = (T_\infty, \ell), \eta)$ be distributed according to $\mu \otimes \mathcal{B}(1/2)$, and let Q_∞ be the image of (θ, η) by the Schaeffer correspondence Φ . The construction of Q_∞ from a tree $\theta \in \mathcal{S}$ allows to specify another class of geodesic rays in Q_∞ , which are defined as follows. These geodesic rays are emanating from the root vertex \emptyset of θ , which can be either e_-^* or e_+^* , depending on the value of η . Consider any infinite path (u_0, u_1, u_2, \dots) in $q = \Phi(\theta, \eta)$ starting from $\emptyset = u_0$, and such that $\ell(u_i) = -i$ for every i . Then necessarily, such a chain is a geodesic ray emanating from \emptyset , because from (3) we have $d_{\text{gr}}^q(u_i, u_j) \geq |i - j|$ for every $i, j \geq 0$, and the other inequality is obviously true.

We call such a geodesic a *proper geodesic ray* emanating from \emptyset . We will see in Corollary 17 below that all geodesic rays emanating from \emptyset are in fact proper.

It should be kept in mind that in the definition of a geodesic ray $(\gamma(i), i \geq 0)$, we can further specify which edge of Q_∞ is used to pass from $\gamma(i)$ to $\gamma(i+1)$. If γ is a proper geodesic, then $\ell(\gamma(i+1)) = \ell(\gamma(i)) - 1$ so this edge has to be an arc drawn in the Schaeffer correspondence between a corner $c^\gamma(i)$ of T_∞ and its successor $\mathcal{S}(c^\gamma(i))$. We will use this several time in the sequel.

The main result of this section shows the existence of cut-points visited by every infinite proper geodesic.




Figure 4: Illustration of the proof of Lemma 7. The tree is represented in solid lines. Every vertex marked by a circled integer corresponds to the last occurrence of this integer along either the left or the right side of the tree. The distinguished geodesic Γ is represented by a thick line.

Theorem 2. *Let $(\theta = (T_\infty, \ell), \eta)$ be distributed according to $\mu \otimes \mathcal{B}(1/2)$, and let Q_∞ be the image of (θ, η) by the Schaeffer correspondence Φ . Almost surely, there exists an infinite sequence of distinct vertices $p_1, p_2, \dots \in V(Q_\infty)$ such that every proper geodesic ray emanating from \emptyset passes through p_1, p_2, \dots*

To prove this theorem we will introduce two specific proper geodesic rays that are in a sense extremal: the minimal and maximal geodesics. We then prove that they meet infinitely often (Lemma 11) and that these meetings points eventually are common to all proper geodesics emanating from \emptyset .

3.2.1 The maximal and minimal geodesics

Recall that if $\theta = (\tau, \ell)$ is a labeled tree in \mathcal{S} with contour sequence $(c_i)_{i \in \mathbb{Z}}$, for every $j \in \mathbb{Z}$ the successor $\mathcal{S}(c_j)$ of c_j is the first corner among c_{j+1}, c_{j+2}, \dots with label $\ell(c_j) - 1$.

Definition 8 (maximal geodesic). Let $\theta = (\tau, \ell) \in \mathcal{S}$. For every corner c of θ , the *maximal geodesic* γ_{\max}^c emanating from c in θ is given by the chain of vertices attached to the iterated successors of c ,


$$\gamma_{\max}^c(i) := \mathcal{V}(\mathcal{S}^{(i)}(c)), \quad i \geq 0,$$

where $\mathcal{S}^{(i)}$ is the i -fold composition of the successor mapping.

Using (3) again, we deduce that the maximal geodesics are indeed geodesic chains in the quadrangulation associated to θ . When $c = c_0$ is the root corner of τ we drop c_0 in the notation

Next, consider only the left part of an infinite labeled tree (τ, ℓ) , which corresponds to the corners (c_0, c_1, c_2, \dots) . We define the *minimal geodesic* γ_{\min} inductively. First, let $\gamma_{\min}(0) = \emptyset$. Suppose that the first n steps $(\gamma_{\min}(0), \dots, \gamma_{\min}(n))$ of γ_{\min} have been constructed. We then set $c^{\gamma_{\min}}(n)$ to be the last corner among c_0, c_1, \dots that is incident to the vertex $\gamma_{\min}(n)$, which implies

One can check by induction that $\ell(\gamma_{\min}(i)) = -i$, thus γ_{\min} is a proper geodesic ray emanating from \emptyset in q . We restrict the definition of the minimal geodesic to the left part of the tree in order to prescribe the behavior of the path when it hits the spine of the tree. Roughly speaking, the minimal geodesic can hit the spine of τ , but it cannot cross it.



The next geometric lemma roughly says that any proper geodesic γ is stuck in between γ_{\min} and γ_{\max} except when γ_{\min} hits the spine of T_∞ in which case γ can visit the right part of the tree T_∞ .

- (i) Suppose that c is a corner incident to $\gamma(i)$ that lies on the left-hand side of τ . Then $c^{\gamma_{\max}}(i) \leq c \leq c^{\gamma_{\min}}(i)$.
- (ii) For every $i \geq 0$, the vertices $\gamma(i+1), \gamma(i+2), \dots$ do not belong to $[\![\emptyset, \gamma(i)]\!]$.
- (iii) For every $i \geq 0$, if $\gamma(i)$ is incident to the left-hand side of τ , then there exists a (unique) $j \leq i$ such that $\gamma_{\min}(j)$ is an ancestor of $\gamma(i)$ in its subtree to the left of the spine. This means that $[\![\emptyset, \gamma(i)]\!]$ contains $\gamma_{\min}(j)$, but $[\![\gamma_{\min}(j), \gamma(i)]\!] \setminus \{\gamma_{\min}(j)\}$ does not intersect the spine of τ .

Next, let γ be a proper geodesic and c be a corner incident to $\gamma(i)$. Since $c^{\gamma_{\max}}(i)$ is the first corner on the left-hand side of τ with label $-i$, and since $\ell(c) = \ell(\gamma(i)) = -i$, if $c \geq c_0$ we must

have $c^{\gamma_{\max}}(i) \leq c$. The inequality $c \leq c^{\gamma_{\min}}(i)$ is true even if $c < c_0$, and is proved by induction. For $i = 0$ it is obvious that any corner incident to $\gamma(0) = \emptyset$ is less than $c^{\gamma_{\min}}(0)$. Suppose that $c \leq c^{\gamma_{\min}}(i)$ for every c incident to $\gamma(i)$. Then this holds in particular for $c = c^{\gamma}(i)$, and we deduce $\mathcal{S}(c^{\gamma}(i)) \leq \mathcal{S}(c^{\gamma_{\min}}(i))$ because \mathcal{S} is non-decreasing when restricted to the (ordered) set of corners of τ with label $-i$. Since $c^{\gamma_{\min}}(i+1)$ is the largest corner incident to $\mathcal{V}(\mathcal{S}(c^{\gamma_{\min}}(i)))$, we obtain that any corner c' incident to $\gamma(i+1)$ satisfies $c' \leq c^{\gamma_{\min}}(i+1)$, unless $\gamma(i+1)$ is a strict ancestor of $\gamma_{\min}(i+1)$ in the subtree to the left of τ that contains $\gamma_{\min}(i+1)$. But by (i), every such ancestor of $\gamma_{\min}(i+1)$ has label at least $-i$, while $\gamma(i+1)$ has label $-i-1$ because γ is a proper geodesic, so the latter obstruction does not occur. This proves (i).

For (ii), since $\ell(\gamma(i)) = -i$, if $\gamma(i+j) \in \llbracket \emptyset, \gamma(i) \rrbracket$ for some $j > 0$, then the cactus bound (4) would imply $d_{\text{gr}}^q(\emptyset, \gamma(i)) \geq 0 - i - 2(-i - j) = i + 2j$, which is impossible since γ is a geodesic and $\emptyset = \gamma(0)$.

We finally prove (iii). It follows from the construction of γ_{\min} that the wanted property is hereditary in i as long as $c^{\gamma}(i) \geq c_0$. In particular, the property holds for $\gamma = \gamma_{\max}$. To show that it is true for any proper geodesic γ , the only problem is when γ visits the right-hand side of τ , which can happen only after a time i where γ hits the spine. So suppose that $\gamma(i)$ belongs to the spine and that $c = c^{\gamma}(i) \leq c_0$. Since γ moves by taking successors, the sequence of corners $c^{\gamma}(j), j \geq i$ can only increase as long as they stay less than c_0 , and after finitely many steps this sequence must leave the right-hand side of τ . In the meantime, it cannot visit the spine, because of (ii). Therefore, at the first time j after i that $\gamma(j)$ leaves the right-hand side of τ , it must make a step from $c^{\gamma}(j-1) \leq 0$ to $\mathcal{S}(c^{\gamma}(j-1)) \geq c_0$. Necessarily, this implies that this successor is also the first corner with label $-j$ on the left-hand side of τ , i.e. $\gamma(j) = \gamma_{\max}(j)$. Being a point of the maximal geodesic, we already noticed that property (iii) holds at this stage, from which we are back to a hereditary situation until the next time k where $c^{\gamma}(k) \leq c_0$, which might happen only at a point where $\gamma_{\min}(k)$ is on the spine. We conclude that the property of (iii) is true at every i where $\gamma(i)$ is incident to the left-hand side, as wanted. \square

Suppose now that for $i \geq 1$ we have $\gamma_{\min}(i) = \gamma_{\max}(i)$. We claim that in fact

$$\mathcal{S}(c^{\gamma_{\max}}(i-1)) = \mathcal{S}(c^{\gamma_{\min}}(i-1)).$$

Indeed by property (i) we have $c^{\gamma_{\max}}(i-1) \leq c^{\gamma_{\min}}(i-1)$, and since these two corners have same label $-i+1$ this implies $\mathcal{S}(c^{\gamma_{\max}}(i-1)) \leq \mathcal{S}(c^{\gamma_{\min}}(i-1))$. The inequality can be strict only if $\gamma_{\min}(i) = \gamma_{\max}(i)$ is an ancestor of $\gamma_{\min}(i-1)$ which is prohibited by property (ii). So if γ is a geodesic such that $c^{\gamma}(i-1)$ lies on the left of T_{∞} then $c^{\gamma_{\max}}(i-1) \leq c^{\gamma}(i-1) \leq c^{\gamma_{\min}}(i-1)$ by property (i), which forces $\mathcal{S}(c^{\gamma_{\max}}(i-1)) = \mathcal{S}(c^{\gamma}(i-1)) = \mathcal{S}(c^{\gamma_{\min}}(i-1))$. In particular $\gamma_{\max}(i) = \gamma_{\min}(i) = \gamma(i)$.

We are going to show that $\gamma_{\max}(i) = \gamma_{\min}(i)$ happens infinitely often, and that any proper geodesic visits only vertices to the left of τ eventually, which will entail Theorem 2.

3.2.2 Inbetween γ_{\min} and γ_{\max}

For $i \geq 0$, denote by A_i the labeled tree consisting of $\gamma_{\min}(i)$ and its descendants in the left side of the tree τ (recall that $\gamma_{\min}(i)$ might be on the spine). Note that this tree may consist only of the single labeled vertex $\gamma_{\min}(i)$. It is clear by construction that A_i is an element of $\mathbf{T}_f^{(-i)}$. Consider the subtree \mathbf{C} of the left-hand side of τ obtained by chopping off the trees $A_i, i \geq 0$, i.e. $v \in \mathbf{C}$ if and only if v is incident to the left-hand side of τ and for every $i \geq 0$, v is not a strict descendent of $\gamma_{\min}(i)$ in the subtree to the left of τ that contains the latter. The tree \mathbf{C} inherits a labeling from ℓ , so it should be seen as an element of \mathcal{S} .

Lemma 10. *Under the law μ , the sequence A_0, A_1, \dots has law $\bigotimes_{i=0}^{\infty} \rho_{-i}$ and is independent of \mathbf{C} .*

Proof of Lemma 10. First of all, remark that A_0 is the tree L_0 grafted to the left-hand side of the first vertex $S(0)$ of the spine of T_∞ . This tree is shortcut by the minimal geodesic γ_{\min} who seeks the successor of $c^{\gamma_{\min}}(0)$ which is the last corner associated to \emptyset . We thus discover the remaining tree $T_\infty \setminus A_0$ step by step in a depth-first search manner by revealing the children of the vertex 1 then the children of 11 (if $11 \in T_\infty$) and so on and so forth in lexicographical order. During this exploration, one can obviously discover the labeling in the same time by sampling at each newly discovered edge a uniform variable $\in \{-1, 0, +1\}$ independently of the past carrying the variation of the label along this edge. We then stop when we discover the first vertex (in lexicographical order) with label -1 , see Fig. 6.




Figure 6: Exploration of T_∞ until the first -1 . The gray trees are unexplored.

This vertex is obviously $\gamma_{\min}(1)$. So far, we thus have explored the part of T_∞ which is composed of the vertices on the left of the segment $[\emptyset, \gamma_{\min}(1)]$ together with all the children of the vertices lying on the ancestral path that link $\gamma_{\min}(1)$ to the spine of T_∞ . These vertices are denoted by $\gamma_{\min}(1) = x_0, x_1, x_2, \dots$ in lexicographical order. Note that some of these vertex can have a label equal to -1 but none of them has a label strictly less than -1 . By standard properties of Galton-Watson tree, the subtrees above x_0, x_1, x_2, \dots are independent of A_0 and of the part of T_∞ explored so far, and form a sequence of independent labeled Galton-Watson trees with laws $\rho_{\ell(x_i)}$. The tree above $x_0 = \gamma_{\min}$ is the tree A_1 that is now shortcut by γ_{\min} who seeks the successor of $c^{\gamma_{\min}}(1)$ (which is the last corner incident to $\gamma_{\min}(1)$). This vertex must lie in the unexplored part of $T_\infty \setminus (A_0 \cup A_1)$. We then continue the exploration of $T_\infty \setminus (A_0 \cup A_1)$ starting with the tree above x_1 in search of the first vertex (for the lexicographical order) with label -2 . The process can be carried out iteratively and yields that A_0, A_1, A_2, \dots has law $\bigotimes_{i \geq 0} \rho_{-i}$ and is independent of $T_\infty \setminus \bigcup_{i \geq 0} A_i$. \square

A key step towards Theorem 2 concerns the intersection points between γ_{\max} and γ_{\min} . Let

$$\mathcal{R} = \{i \geq 0 : \gamma_{\max}(i) = \gamma_{\min}(i)\},$$

and note that the set $\{\gamma_{\max}(i) : i \in \mathcal{R}\}$ is equal to the intersection of the images of $\gamma_{\max}, \gamma_{\min}$. Indeed, from the fact that $\gamma_{\max}, \gamma_{\min}$ are both geodesics started from \emptyset , $\gamma_{\max}(i) = \gamma_{\min}(j)$ automatically implies $i = j$. Recall that a (discrete) regenerative set is a random set of the form $\mathcal{G} = \{G_0 + G_1 + \dots + G_n, n \geq 0\}$, where G_1, G_2, \dots are i.i.d. random variables with values in \mathbb{N} , independent of G_0 . It is called aperiodic if the greatest common divisor of the support of the




Figure 7: The second step of the exploration.

law of G_1 is 1. If \mathcal{G} is a regenerative set, then by the renewal theorem its asymptotic frequency exists and is given by

$$|\mathcal{G}| = \lim_{n \geq 1} \frac{\#\mathcal{G} \cap \{1, \dots, n\}}{n} = \frac{1}{\mathbb{E}[G_1]},$$

and if \mathcal{G} is aperiodic then $|\mathcal{G}| = \lim_{i \rightarrow \infty} \mathbb{P}(i \in \mathcal{G})$

Lemma 11. *The set \mathcal{R} is a (discrete) regenerative set, and $|\mathcal{R}| > 0$ a.s.*

Proof. We first note that the maximal geodesic only visits vertices in the trees $A_i, i \geq 0$, by (iii) in Lemma 9. More precisely, for every $i \geq 0$, $\gamma_{\max}(i)$ belongs to the tree $A_{j(i)}$ where $j(i)$ is the first index $j \leq i$ such that $\min\{\ell(v) : v \in A_j\} = -i$. Here we make a slight abuse of language by viewing A_i as a (labeled) subgraph of (T_∞, ℓ) rather than a tree in its own right.

In particular, we deduce that $\gamma_{\max}(i) = \gamma_{\min}(i)$ if and only if $j(i) = i$, that is $\inf\{\ell(v) : v \in A_0 \cup \dots \cup A_{i-1}\} = -i + 1$. Otherwise said, letting $\Delta_i = \max\{-i - \ell(v) : v \in A_i\} \geq 0$, then $\max_{0 \leq j \leq i-1} (j + \Delta_j) = i - 1$, and this gives

$$\mathcal{R} = \mathbb{Z}_+ \setminus \bigcup_{j \geq 0} (j, j + \Delta_j].$$

Yet otherwise said, \mathcal{R} has same distribution as the set $\{G_0 + G_1 + \dots + G_n : n \geq 0\}$ where $G_0 = 0$ and the variables G_1, G_2, \dots are i.i.d. with $G_1 = \inf\{i > 0 : \max\{j + \Delta_j : 0 \leq j < i\} < i\}$. Now, by Lemma 10, the random variables $(\Delta_i, i \geq 0)$ are independent and distributed as $\max\{-\ell(v) : v \in \tau\}$ where (τ, ℓ) has law ρ_0 . By symmetry, this is the same as the law of $\max\{\ell(v) : v \in \tau\}$, still under ρ_0 . We will use the following lemma whose proof is postponed to Section 4:

Lemma 12. *It holds that, as $m \rightarrow \infty$,*

$$\mathbb{P}(\Delta_0 \geq m) \sim \frac{2}{m^2},$$

the symbol \sim meaning that the quotient of both sides tends to 1.

We note that

$$\begin{aligned}
\mathbb{P}(i \in \mathcal{R}) &= \mathbb{P}(\max\{j + \Delta_j : 0 \leq j \leq i-1\} < i) \\
&= \prod_{j=0}^{i-1} (1 - \mathbb{P}(\Delta_0 \geq i-j)) \\
&= \prod_{j=1}^i (1 - \mathbb{P}(\Delta_0 \geq j)),
\end{aligned}$$

which as $i \rightarrow \infty$ decreases to a strictly positive limit by Lemma 12 and the (obvious) fact that $\mathbb{P}(\Delta_0 = 0) > 0$. This concludes the proof of Lemma 11. \square

Note that if \mathcal{G} is a given infinite subset of \mathbb{Z}_+ , then Lemma 11 shows that $\mathcal{R} \cap \mathcal{G}$ is infinite almost surely, because

$$\mathbb{P}(\#\mathcal{R} \cap \mathcal{G} = \infty) \geq \lim_{i \rightarrow \infty, i \in \mathcal{G}} \mathbb{P}(i \in \mathcal{R}) > 0,$$

so the probability that $\#\mathcal{R} \cap \mathcal{G} = \infty$ has to be 1 by the Hewitt-Savage 0-1 law.

3.2.3 Leaving the spine

Our next step towards Theorem 2 is to prove that γ_{\max} and γ_{\min} eventually leave the spine for ever. We begin with the case of the maximal geodesic. Let

$$\mathcal{R}' = \{i \geq 0 : \gamma_{\max}(i) \in \{S(j), j \geq 0\}\}$$

be the set of times where γ_{\max} hits the spine.

Lemma 13. *Almost surely, the set \mathcal{R}' is finite.*

Proof. Let L_0, L_1, L_2, \dots be the subtrees to the left of the spine of T_∞ . Recall the notation $\ell(S(i)) = X_i$. Then note that the i -th vertex $S(i)$ of the spine is on γ_{\max} if and only if

$$\min\{\ell(v) : v \in L_j\} > \ell(S(i)), \quad \text{for every } j \in \{0, 1, \dots, i-1\}.$$

Now, for $i \geq 0$ let $\sigma_i = \inf\{n \geq 0 : X_n = -i\}$, and let L'_i be the forest made of the trees L_{σ_i+j} , $0 \leq j < \sigma_{i+1} - \sigma_i$. Then, if we let

$$\Delta'_i = \max_{v \in L'_i} -\ell(v) - i, \quad i \geq 0,$$

then we obtain that

$$\mathcal{R}' = \mathbb{Z}_+ \setminus \bigcup_{j \geq 0} (j, j + \Delta'_j].$$

Furthermore the Δ'_i are independent and identically distributed. At this point the situation is similar to the setting of Lemma 11 and we also rely on an external lemma whose proof is postponed to Section 4:

Lemma 14. *As $m \rightarrow \infty$, it holds that*

$$\mathbb{P}(\Delta'_0 \geq m) \sim \frac{2}{m}.$$

By the very same argument as for the computation of $\mathbb{P}(i \in \mathcal{R})$ we have using Lemma 14

$$\begin{aligned}
\mathbb{P}(i \in \mathcal{R}') &= \prod_{j=1}^i (1 - \mathbb{P}(\Delta'_0 \geq j)) \\
&= \exp \left(- \sum_{j=1}^i -\log(1 - \mathbb{P}(\Delta'_0 \geq j)) \right) \\
&= \exp \left(- 2 \log m(1 + o(1)) \right) \\
&= m^{-2+o(1)}.
\end{aligned}$$

In particular, $\mathbb{P}(i \in \mathcal{R}')$ is summable, so that, by the Borel-Cantelli Lemma, $i \notin \mathcal{R}'$ for every i large enough, as wanted. \square

We deduce the following key property of the minimal geodesic.

Proposition 15. *Almost-surely, the minimal geodesic γ_{\min} hits the spine a finite number of times:*

$$\#\{i \geq 0 : \gamma_{\min}(i) \in \{S(j), j \geq 0\}\} < \infty.$$

Proof. Assume by contradiction that γ_{\min} hits the spine infinitely often with positive probability, and let \mathcal{G} be the set of such intersections. Then \mathcal{G} is measurable with respect to the subtree \mathbb{C} obtained by chopping the trees A_0, A_1, A_2, \dots off θ . By Lemma 10, we obtain that \mathcal{G} is independent of $(A_i, i \geq 0)$, whence \mathcal{G} is independent of the set \mathcal{R} of intersections times of γ_{\max} with γ_{\min} . Conditionally given \mathcal{G} , in restriction to the event that the latter is infinite, we then conclude from the discussion after Lemma 11 that $\mathcal{R} \cap \mathcal{G}$ is infinite almost-surely. But this is in contradiction with Lemma 13. \square

3.2.4 Proof of Theorem 2

Proof of Theorem 2. Let

$$i_* = \max \{j \geq 0 : \gamma_{\min}(j) \in \{S(0), S(1), \dots\}\}$$

be the last time when γ_{\min} hits the spine, and

$$c_* = \min \{c : \mathcal{V}(c) = \gamma_{\min}(i_*)\},$$

the minimal corner incident to $\gamma_{\min}(i_*)$. Note that $c_* \leq c_0$, that is, c_* lies on the right-hand side of T_∞ . Let γ be a proper geodesic emanating from \emptyset . Recall from the proof of (iii) in Lemma 9 that $\gamma(i)$ belongs to the spine if and only if $\gamma(i) = \gamma_{\min}(i)$ and $\gamma_{\min}(i)$ belongs to the spine. Moreover, after each visit to the spine, the corners used by γ increase until the next visit to the spine. It follows that $c^\gamma(i) \geq c_*$ for every $i \geq 0$. In particular, $\gamma(i)$ is incident to the left-hand side of T_∞ for every $i > i_0$, where $i_0 = -\min\{\ell(c) : c_* \leq c \leq c_0\}$ (the latter depends only on (T_∞, ℓ) and not on the choice of γ).

Now by (i) in Lemma 9, we deduce that $c^{\gamma_{\max}}(i) \leq c^\gamma(i) \leq c^{\gamma_{\min}}(i)$ for every $i > i_0$. In particular, for every $i > i_0 + 1$ such that $i \in \mathcal{R}$, as defined around Lemma 11, we have $\gamma(i) = \gamma_{\max}(i) = \gamma_{\min}(i)$ by the discussion after the proof of Lemma 9. Letting p_1, p_2, \dots be the ordered list of points of $\{\gamma_{\max}(i) : i \in \mathcal{R}, i > i_0\}$, which is infinite by Lemma 11, we thus see that every proper geodesic γ has to visit the points p_1, p_2, \dots . This concludes the proof of Theorem 2. \square




Figure 8: Illustration of the proof of Theorem 2: A proper geodesic can “swirl” around T_∞ by traversing the spine, but is eventually trapped between the minimal and maximal geodesics

3.3 End of the proof of Theorem 1

Lemma 16. *Almost surely, the function $z \mapsto d_{\text{gr}}^{Q_\infty}(z, e_-^*) - d_{\text{gr}}^{Q_\infty}(z, e_+^*)$ from $V(Q_\infty)$ to $\{-1, 1\}$ is almost constant., i.e. is constant except for finitely many $z \in Q_\infty$.*

Proof. This statement is a property of the UIPQ, but for the purposes of the proof, we will assume that Q_∞ is constructed from a tree $\theta = (T_\infty, \ell)$ with law μ and an independent parameter η with $\mathcal{B}(1/2)$ distribution, by applying the Schaeffer correspondence Φ . This allows to specify the class of proper geodesic rays among all geodesic rays emanating from \emptyset .

First, let us assume that $\emptyset = e_-^*$, meaning that $\eta = 0$. Let γ_{\max} be the maximal geodesic so that $\gamma_{\max}(0) = \emptyset = e_-^*$, and $\gamma_{\max}(1) = e_+^*$. It is also a proper geodesic ray, so that $\ell(\gamma_{\max}(i)) = -i$ for every $i \geq 0$.

Note that if γ is a geodesic from \emptyset to $\gamma_{\max}(i)$ for some $i \geq 0$, then necessarily $\ell(\gamma(j)) = -j$ for every $j \in \{0, 1, 2, \dots, i\}$, the reason being that the labels of two neighboring vertices in Q_∞ differ by at most 1.

Now let Γ be the distinguished geodesic ray which starts from $e_-^* = \emptyset$ constructed from Q_∞ by first recovering the Chassaing-Durhuus tree $(\tau, \ell) = \Phi^{-1}(Q_\infty)$ and then constructing Γ as we did just before Lemma 7, and let $R \geq 0$. Applying Lemma 7, we obtain the existence of $R' \geq R$ such that the vertex $\gamma_{\max}(R' + 1)$, which does not belong to $B_{\mathbf{Q}, R'}(Q_\infty)$, can be linked to \emptyset by a geodesic γ such that $\gamma(i) = \Gamma(i)$ for $i \in \{0, 1, \dots, R\}$. Since $\ell(\gamma(R' + 1)) = \ell(\gamma_{\max}(R' + 1)) = -(R' + 1)$, we deduce from the above discussion that $\ell(\gamma(i)) = -i$ for every $i \in \{0, 1, \dots, R'\}$, so in particular, $\ell(\Gamma(i)) = -i$ for every $i \in \{0, 1, \dots, R\}$. Since R was arbitrary, we deduce that the distinguished geodesic Γ is proper.

By Theorem 2, we get that Γ and γ_{\max} meet infinitely often. In particular, for every $\alpha \in (0, 1)$, we can find $R = R(\alpha)$ such that with probability at least $1 - \alpha$, there exists $I \in \{1, 2, \dots, R\}$ such that $\Gamma(I) = \gamma_{\max}(I)$. From now on we argue on this event. Applying Lemma 7 again, we can find R' such that for every $z \in V(Q_\infty) \setminus B_{\mathbf{Q}, R'}(Q_\infty)$, one can link \emptyset to z by a geodesic γ whose R' first steps coincide with those of Γ . But since $\Gamma(I) = \gamma_{\max}(I)$, we can replace the first I steps of γ by those of γ_{\max} , and obtain a new geodesic from \emptyset to z , whose first step goes from e_-^* to e_+^* . Since this holds for any z at distance at least $R' + 1$ from

e_-^* , we obtain that $d_{\text{gr}}^{Q_\infty}(z, e_-^*) - d_{\text{gr}}^{Q_\infty}(z, e_+^*) = 1$ for every z at distance at least $R' + 1$ from e_-^* . Since α was arbitrary, we obtain the desired result in the case $\eta = 0$.

To treat the case $\eta = 1$, we use the obvious fact that if \overleftarrow{Q}_∞ is the same quadrangulation as Q_∞ , but where the root edge has the reverse orientation, then \overleftarrow{Q}_∞ has the same distribution as Q_∞ . Moreover, $\overleftarrow{Q}_\infty = \Phi(\theta, 1 - \eta)$ so on the event $\{\eta = 1\}$ we are back to the situation $\eta = 0$ by arguing on \overleftarrow{Q}_∞ instead of Q_∞ . \square

From this, it is easy to prove (6), which will complete the proof of Theorem 1. Indeed, if x and y are neighboring vertices in Q_∞ we can pick an edge e such that $e_- = x$ and $e_+ = y$. By Proposition 19 below, the quadrangulation $Q_\infty^{(e)}$ re-rooted at e has the same almost sure properties as Q_∞ . In particular, almost surely the function $z \mapsto d(x, z) - d(y, z)$ is almost constant. But by reasoning on every step of a chain from x to y , the same holds for any $x, y \in Q_\infty$. This constant has to be $\ell(x) - \ell(y)$. Indeed let us consider γ_x and γ_y two maximal geodesics emanating from a corner associated to x resp. y . From properties of the Schaeffer construction, these two geodesics merge at some vertex $\gamma_x(i) = \gamma_y(i + \ell(y) - \ell(x))$ for some $i \in \{0, 1, 2, \dots\}$, and $\gamma_x(j) = \gamma_y(j + \ell(y) - \ell(x))$ for every $j \geq i$. Hence

$$\lim_{z \rightarrow \infty} \left(d_{\text{gr}}^{Q_\infty}(x, z) - d_{\text{gr}}^{Q_\infty}(y, z) \right) = \lim_{\substack{z \rightarrow \infty \\ z \in \gamma_x \cap \gamma_y}} \left(d_{\text{gr}}^{Q_\infty}(x, z) - d_{\text{gr}}^{Q_\infty}(y, z) \right) = \ell(x) - \ell(y).$$

Corollary 17. *Every geodesic ray emanating from \emptyset is proper.*

Proof. Let γ be a geodesic ray and let $i_0 \geq 1$ fixed. Applying (7) we get

$$\begin{aligned} \ell(\gamma(i_0)) &= \lim_{z \rightarrow \infty} \left(d_{\text{gr}}^{Q_\infty}(\gamma(i_0), z) - d_{\text{gr}}^{Q_\infty}(\emptyset, z) \right) \\ &= \lim_{i \rightarrow \infty} \left(d_{\text{gr}}^{Q_\infty}(\gamma(i_0), \gamma(i)) - d_{\text{gr}}^{Q_\infty}(\emptyset, \gamma(i)) \right). \end{aligned}$$

On the other hand, since γ is a geodesic, for $i \geq i_0$ we have $d_{\text{gr}}^{Q_\infty}(\gamma(i_0), \gamma(i)) = i - i_0$, which implies that $\ell(\gamma(i_0)) = -i_0$. This allows to conclude since i_0 was arbitrary. \square

4 Scaling limits for (T_∞, ℓ)

This section is devoted to the study of the scaling limits of the contour functions describing the tree (T_∞, ℓ) . It also contains the proof of Lemmas 12 and 14 used during the proof of Theorem 2.

4.1 Contour functions

Coding of a tree. Let us recall a useful encoding of labeled trees by functions. A finite labeled tree $\theta = (\tau, \ell)$ can be encoded by a pair (C_θ, V_θ) , where $C_\theta = (C_\theta(t))_{0 \leq t \leq 2|\theta|}$ is the contour function of τ and $V_\theta = (V_\theta(t))_{0 \leq t \leq 2|\theta|}$ is the labeling contour function of θ . To define these contour functions, we let $(c_0, c_1, \dots, c_{2|\theta|-1})$ be the contour sequence of corners of τ . Then $C_\theta(i)$ is the distance of $\mathcal{V}(c_i)$ to the root \emptyset in τ for $0 \leq i \leq 2|\theta| - 1$, and we let $C_\theta(2|\theta|) = 0$. Furthermore, we let $V_\theta(i) = \ell(c_i)$ for $0 \leq i \leq 2|\theta| - 1$, and then $V_\theta(2|\theta|) = \ell(c_0)$. Finally, we extend C_θ, V_θ to continuous functions on $[0, 2|\theta|]$ by linear interpolation between integer times (we will generally ignore these interpolations in what follows). See Figure 9 for an example. A finite labeled tree is uniquely determined by its pair of contour functions.

If $\theta = (\tau, \ell)$ has law ρ_l for some $l \in \mathbb{Z}$, then it is easy and well-known that C_θ is a simple random walk, stopped at time $\kappa_1 - 1$, where κ_1 the first hitting time of -1 . For this reason, the process C_θ is sometimes extended to $[0, 2|\theta| + 1]$ by taking a final step of amplitude -1 , this will help explain the construction of the process C below.




Figure 9: A labeled tree θ and its pair of contour functions (C_θ, V_θ) .

Coding of a forest. If we now consider a sequence of trees $\theta_i = (\tau_i, \ell_i), i \geq 0$ respectively in $\mathbf{T}_f^{(\ell_i)}$, then we can concatenate the (extended) processes C_{θ_i} in a process

$$C(k) = C_{\theta_i} \left(k - \sum_{j < i} (2|\theta_j| + 1) \right) - i, \quad \sum_{j < i} (2|\theta_j| + 1) \leq k < \sum_{j \leq i} (2|\theta_j| + 1)$$

We view C as the contour function of an infinite forest $(\theta_0, \theta_1, \dots)$, in which C takes a -1 step at every newly visited tree. If we let $\underline{C}(i) = \inf_{0 \leq j \leq i} C_j$, then the process C_{θ_i} can be recovered as

$$C_{\theta_i}(j) = i + C(\kappa_i + j) = C(\kappa_i + j) - \underline{C}(\kappa_i + j), \quad 0 \leq j < \kappa_{i+1} - \kappa_i = 2|\theta_i| + 1,$$

where $\kappa_i = \inf\{n \geq 0 : C(n) = -i\}$. We could also have chosen to take a 0 or $+1$ steps at every newly visited tree: these contour functions are respectively given by $C - \underline{C}$ and $C - 2\underline{C}$. We will see that it is easier to deal with C but $C - 2\underline{C}$ also plays a natural role, see the next paragraph.




Figure 10: Different codings of a forest

As for the labeling function V , it is defined accordingly by concatenation of the processes $V_{\theta_i}, i \geq 0$, so for $i \geq 0$,

$$V(k + \kappa_i) = V_{\theta_i}(k), \quad 0 \leq k < \kappa_{i+1} - \kappa_i.$$

4.2 Scaling limits of contour functions

To describe the left part of the tree T_∞ one thus would like to understand the contour functions associated to the sequence of trees L_0, L_1, L_2, \dots . It turns out that it is easier to first deal with the contour and label processes of the labeled forest L_0, L_1, \dots , in which we subtracted the label X_i of the root of L_i to the labels of all vertices of L_i . These sequences are denoted by (C, V) . Indeed, after doing this operation, the relabeled trees we obtain form an i.i.d. sequence with law ρ_0 . In particular the process C is a standard simple random walk with unit jumps. The “true” label $\ell(\mathcal{V}(c_n))$ of the n -th vertex visited in the contour order of T_∞ is then given by the formula

$$V'(n) = X_{-\underline{C}(n)} + V(n), \quad (10)$$

because $-\underline{C}(n)$ is the index of the tree to which $\mathcal{V}(c_n)$ belongs. Obviously, the contour function C is unchanged by this operation. Note that the labels on the spine X is independent of (C, V) . We will introduce the scaling limits of X and (C, V) separately. First, the Donsker invariance principle implies that

$$\left(\frac{1}{m} \sqrt{\frac{3}{2}} X_{m^2 t} \right)_{t \geq 0} \xrightarrow[m \rightarrow \infty]{(d)} (\beta_t)_{t \geq 0}, \quad (11)$$

where β is a standard Brownian motion. On the other hand, let $(B_s, s \geq 0)$ be a standard Brownian motion, and $\underline{B}_s = \inf_{0 \leq u \leq s} B_u$ be its infimum process. For $s \geq 0$, we let $\hat{B}_s = B_s - \underline{B}_s$, so that $\hat{B} = (\hat{B}_s, s \geq 0)$ is a standard reflected Brownian motion with local time at zero given by $(-\underline{B}_s, s \geq 0)$, by a famous theorem due to Lévy. Now, conditionally given \hat{B} , let $(Z_s, s \geq 0)$ be a centered Gaussian process whose covariance function is given by

$$E[(Z_s - Z_t)^2] = \hat{B}_s + \hat{B}_t - 2 \inf_{u \in [s \wedge t, s \vee t]} \hat{B}_u.$$

The process Z has a continuous modification [24] and this is the one we will deal with henceforth. The pair (\hat{B}, Z) is called the Brownian snake (or sometimes the head of the Brownian snake) driven by the reflected Brownian motion \hat{B} . The reader can refer to the monograph [24] for a detailed account of the Brownian snake. Then we have the joint convergence in distribution for the uniform norm over every compact interval:

$$\left(\frac{1}{m^2} C(m^4 t), \frac{1}{m^2} \underline{C}(m^4 t), \frac{1}{m} \sqrt{\frac{3}{2}} V(m^4 t) \right)_{t \geq 0} \xrightarrow[n \rightarrow \infty]{(d)} (B_t, \underline{B}_t, Z_t)_{t \geq 0}, \quad (12)$$

this convergence holds jointly with (11) and (β_t) is independent of the triplet $(B_t, \underline{B}_t, Z_t)$. See for instance [31, Theorem 3] for a similar statement, from which the present one can be deduced easily. One difference is that [31] deals with the so-called *height process* of the trees rather than the contour process, but the convergence of the latter is indeed a consequence of the convergence of height processes, as discussed in [16, Section 2.4]. Convergences (11) and (12) entail that the contour functions (C, V') of the forest L_0, L_1, \dots of the left part of T_∞ admit the following scaling limit

$$\left(\frac{1}{m^2} C(m^4 t), \frac{1}{m} \sqrt{\frac{3}{2}} V'(m^4 t) \right)_{t \geq 0} \xrightarrow[m \rightarrow \infty]{(d)} (B_t, \beta_{-\underline{B}_t} + Z_t)_{t \geq 0}. \quad (13)$$

To deal with the right part of T_∞ , we just remark that if \tilde{C}, \tilde{V} and \tilde{V}' are defined from the forest R_0, R_1, \dots in the same way as C, V and V' were defined from L_0, L_1, \dots then we have the analogous of (10)

$$\tilde{V}'(n) = X_{-\tilde{C}(n)} + \tilde{V}(n),$$

where X is still the random walk of the labels on the spine. Remark that (C, V) and (\tilde{C}, \tilde{V}) are independent and identically distributed and also independent of X . Hence the convergence (12) also holds for (\tilde{C}, \tilde{V}) namely jointly with (11) and (12) we have the convergence in distribution

$$\left(\frac{1}{m^2} \tilde{C}(m^4 t), \frac{1}{m^2} \tilde{\underline{C}}(m^4 t), \frac{1}{m} \sqrt{\frac{3}{2}} \tilde{V}(m^4 t) \right)_{t \geq 0} \xrightarrow[n \rightarrow \infty]{(d)} (\tilde{B}_t, \tilde{\underline{B}}_t, \tilde{Z}_t)_{t \geq 0}, \quad (14)$$

where $(\tilde{B}, \tilde{\underline{B}}, \tilde{Z})$ is an independent copy of (B, \underline{B}, Z) also independent of β . Finally, the scaling limits of \tilde{C} and \tilde{V}' is given by an analogous formula as (13) after replacing B, \underline{B} and Z by their tilde-analogs.

4.3 The continuous tree $(\mathcal{T}_\infty, \mathcal{Z})$

Let us give a slightly different point of view on these scaling limit results for the contour functions (C, V) and (\tilde{C}, \tilde{V}) . The results of this paragraph will not be used in the sequel, we thus leave the details to the reader.

First of all, we remark that by a famous theorem of Pitman (see [33, Chapter VI]) the processes $B - 2\underline{B}$ and $\tilde{B} - 2\underline{\tilde{B}}$ are two independent three-dimensional Bessel processes. These two processes thus give the scaling limits of the two contour functions of the left and right part of T_∞ in which we make a $+1$ steps at the end of every visited tree. We now let

$$\mathfrak{R}(t) = \begin{cases} B_t - 2\underline{B}_t & \text{if } t \geq 0, \\ \tilde{B}_{-t} - 2\underline{\tilde{B}}_{-t} & \text{if } t \leq 0, \end{cases}$$

and for every $s, t \in \mathbb{R}$, we define

$$\overline{st} = \begin{cases} [s \wedge t, s \vee t] & \text{if } st \geq 0, \\ (-\infty, s \wedge t] \cup [s \vee t, \infty) & \text{if } st < 0. \end{cases}$$

Finally we define a pseudo-distance on \mathbb{R} by letting

$$d_{\mathfrak{R}}(s, t) = \mathfrak{R}_s + \mathfrak{R}_t - 2 \inf_{r \in \overline{st}} \mathfrak{R}_r.$$

The quotient space $\mathcal{T}_\infty = \mathbb{R}/(d_{\mathfrak{R}} = 0)$ equipped with the quotient distance $d_{\mathfrak{R}}$ is an infinite tree that is the scaling limit of the tree T_∞ , that is $\lambda \cdot T_\infty \rightarrow \mathcal{T}_\infty$ in distribution for the Gromov-Hausdorff distance [10] as $\lambda \rightarrow 0$.

Furthermore, conditionally on \mathfrak{R} , we consider a real-valued centered Gaussian process \mathcal{Z}_t indexed by \mathbb{R} whose covariance function is prescribed by

$$E[(\mathcal{Z}_s - \mathcal{Z}_t)^2] = d_{\mathfrak{R}}(s, t).$$

Similarly to the case of Z , the process \mathcal{Z} has a continuous modification that we consider from now on. In words, the process \mathcal{Z} can be interpreted as the Brownian motion indexed by the infinite tree \mathcal{T}_∞ . We claim that conditionally on C and on \tilde{C} we have the equality in distribution

$$\left(\mathbf{1}_{t \geq 0} (Z_t + \beta_{-\underline{B}_t}) + \mathbf{1}_{t \leq 0} (\tilde{Z}_{-t} + \beta_{-\underline{\tilde{B}}_{-t}}) \right) \stackrel{(d)}{=} (\mathcal{Z}_t)_{t \in \mathbb{R}},$$

which can be check by looking at the covariance functions of these Gaussian processes. To conclude, this interpretation enables us to fully understand that the labeled tree T_∞ converges, in the scaling limit, towards a non-compact random tree \mathcal{T}_∞ encoded by a pair of independent Bessel processes with the Brownian labeling \mathcal{Z} . This object should play a crucial role to describe the scaling limit of the UIPQ in the Gromov-Hausdorff sense. We plan to study this in future work.

4.4 Proofs of Lemmas 12 and 14

In this section we proceed to the proof of the lemmas used during the proof of Theorem 2. Although these lemmas seem combinatorial in nature, the constant appearing in the equivalents are obtained by using the continuous scaling limits of the contour processes of the tree T_∞ .

Recall that Δ_0 is distributed as $\max\{\ell(v) : v \in \tau\}$ under ρ_0 .

Lemma 12. *It holds that, as $m \rightarrow \infty$,*

$$\mathbb{P}(\Delta_0 \geq m) \sim \frac{2}{m^2}.$$

Proof. Consider an infinite sequence $(\theta_i, \ell_i)_{i \geq 0}$ of independent labeled trees with law ρ_0 . We let (C, V) be the contour and label sequences of this forest, as defined in the last section. Recall the notation $\kappa_i = \inf\{n \geq 0 : C(n) = -i\}$. The convergence (12), together with standard arguments relying on the fact that for a given level $-x$, the first hitting time $T_x = \inf\{t \geq 0 : B_t < -x\}$ (the definition is chosen so that T is the right-continuous inverse of the function $-\underline{B}$) is almost surely not a local minimum for the function B , entails that

$$\frac{1}{m} \sqrt{\frac{3}{2}} \max\{V(i) : 0 \leq i \leq \kappa_{m^2}\} \xrightarrow[m \rightarrow \infty]{(d)} \sup\{Z_t : 0 \leq t \leq T_1\}. \quad (15)$$

By excursion theory for the Brownian snake [24], it holds that

$$Z^{(a)} = (Z_{T_{a-}+s}, 0 \leq s \leq T_a - T_{a-}), \quad a \geq 0$$

is a Poisson point process with intensity $2\mathbb{N}_0(dZ)$, where \mathbb{N}_0 is the so-called Itô excursion measure of the Brownian snake, and with standard Brownian spatial displacements. In particular, it is known, by [28, Lemma 2.1] and invariance in distribution of the process Z under reflection, that

$$2\mathbb{N}_0(\sup Z > x) = \frac{3}{x^2}, \quad x > 0.$$

Therefore, by standard properties of Poisson random measures,

$$\mathbb{P}(\sup\{Z_t : 0 \leq t \leq T_1\} \leq x) = \mathbb{P}(\sup\{\sup Z^{(a)} : 0 \leq a \leq 1\} \leq x) = \exp(-3/x^2).$$

Moreover, since the paths $(V(\kappa_i + j), 0 \leq j \leq \kappa_{i+1} - \kappa_i), i \geq 0$ are i.i.d. and $\max\{V(j) : 0 \leq j \leq \kappa_1\}$ has same law as Δ_0 , we obtain that for every $x > 0$,

$$\begin{aligned} \left(1 - \mathbb{P}(\Delta_0 > mx\sqrt{\frac{2}{3}})\right)^{m^2+1} &= \mathbb{P}\left(\frac{1}{m}\sqrt{\frac{3}{2}} \max\{V(i) : 0 \leq i \leq \kappa_{m^2}\} \leq x\right) \\ &\xrightarrow[m \rightarrow \infty]{} \exp(-3/x^2), \end{aligned}$$

where in the last step we used (15) and the fact that the random variable $\sup\{Z_t : 0 \leq t \leq T_1\}$ has a diffuse law. This entails

$$m^2 \mathbb{P}(\Delta_0 > m) \rightarrow 2,$$

concluding the proof of Lemma 12. □

Next, using the notation of Section 4.2, we let

$$\kappa'_i = \inf\{j \geq 0 : X_{-\underline{C}(j)} = -i\}.$$

We recall the definition of the sequence $(\Delta'_i, i \geq 0)$ introduced in Lemma 13

$$\Delta'_i = \max\{-V'(\kappa'_i + j), 0 \leq j \leq \kappa'_{i+1} - \kappa'_i\} - i.$$

Lemma 14. *As $m \rightarrow \infty$, it holds that*

$$\mathbb{P}(\Delta'_0 \geq m) \sim \frac{2}{m}.$$

Proof. The sequence Δ'_i admits the alternative representation

$$\Delta'_i = \max\{-V(\kappa'_i + j) - X_{-\underline{C}(\kappa'_i + j)} + \underline{X}_{-\underline{C}(\kappa'_i + j)}, 0 \leq j \leq \kappa'_{i+1} - \kappa'_i\}, \quad i \geq 0, \quad (16)$$

where $\underline{X}_i = \min\{X_j, 0 \leq j \leq i\}$, so that

$$\max_{0 \leq i \leq m} \Delta'_i = \max\{-V(j) - X_{-\underline{C}(j)} + \underline{X}_{-\underline{C}(j)} : 0 \leq j \leq \kappa'_m\}.$$

Note also that $(\Delta'_i, i \geq 0)$ is an i.i.d. sequence. This can be proved by exploring the sequence L_0, L_1, \dots in a Markovian way, in the spirit of the proof of Lemma 10, and we leave this fact to the reader. Convergences (11) and (12) entail that

$$\left(\frac{1}{m}\sqrt{\frac{3}{2}}(V(m^4t) + X_{-\underline{C}(m^4t)} - \underline{X}_{-\underline{C}(m^4t)})\right)_{t \geq 0} \xrightarrow{m \rightarrow \infty} (Z_t + \hat{\beta}_{-\underline{B}_t} t \geq 0),$$

where $\hat{\beta}_x = \beta_x - \underline{\beta}_x$. Let $T'_x = \inf\{t \geq 0 : \beta_{-\underline{B}_t} < -x\} = T_{\tau_x}$, where $\tau_x = \inf\{u \geq 0 : \beta_u < -x\}$ is the right-continuous inverse of $-\underline{\beta}$. For a given $x \geq 0$, it is easy to check that the time T'_x is not a local minimum for the process $(\beta_{-\underline{B}_t}, t \geq 0)$. Therefore, the previous convergence entails

$$\left(\frac{1}{m}\sqrt{\frac{3}{2}}(V'(m^4t) - \underline{X}_{-\underline{C}(m^4t)})\right)_{0 \leq t \leq \kappa'_m/m^4} \xrightarrow{m \rightarrow \infty} (Z_t + \hat{\beta}_{-\underline{B}_t})_{0 \leq t \leq T'_{\sqrt{3/2}}} \quad (17)$$

Let us fix $x \geq 0$. We claim that

$$\mathbb{P}\left(\sup\{Z_t - \hat{\beta}_{-\underline{B}_t} : 0 \leq t \leq T'_{\sqrt{3/2}}\} \leq x\right) = \exp(-\sqrt{6}/x). \quad (18)$$

To show this, recall the notation $Z^{(a)}$ used in the proof of Lemma 12. Then,

$$\begin{aligned} & \sup\{Z_t - \hat{\beta}_{-\underline{B}_t} : 0 \leq t \leq T'_{\sqrt{3/2}}\} \\ &= \sup\{\sup\{Z_{T_{a-}+s} : 0 \leq s \leq T_a - T_{a-}\} - \hat{\beta}_a : 0 \leq a \leq \tau_{\sqrt{3/2}}\} \\ &= \sup\{\sup Z^{(a)} - \hat{\beta}_a : 0 \leq a \leq \tau_{\sqrt{3/2}}\}. \end{aligned}$$

Since $(Z^{(a)}, a \geq 0)$ is a homogeneous Poisson process, standard properties of Poisson measures entail that conditionally given β ,

$$\begin{aligned} \mathbb{P}\left(\sup\{Z_t - \hat{\beta}_{-\underline{B}_t} : 0 \leq t \leq T'_{\sqrt{3/2}}\} \leq x \mid \beta\right) &= \exp\left(-\int_0^{\tau_{\sqrt{3/2}}} da \, 2\mathbb{N}_0(\sup Z - \hat{\beta}_a \geq x)\right) \\ &= \exp\left(-\int_0^{\tau_{\sqrt{3/2}}} \frac{da}{(x + \hat{\beta}_a)^2}\right) \\ &= \exp\left(-\sum_{0 < y \leq \sqrt{3/2}} \int_0^{\tau_y - \tau_{y-}} \frac{da}{(x + \beta_a^{(y)})^2}\right) \end{aligned}$$

where

$$\beta_a^{(y)} = \hat{\beta}_{\tau_y + a} = y + \beta_{\tau_y + a}, \quad 0 \leq a \leq \tau_y - \tau_{y-}$$

is the excursion of β above its minimum at level $-y$. Taking expectations and using Itô's excursion theory, we obtain

$$\mathbb{P}\left(\sup\{Z_t - \hat{\beta}_{-\underline{B}_t} : 0 \leq t \leq T'_{\sqrt{3/2}}\} \leq x\right) = \exp\left(-\sqrt{\frac{3}{2}} \int 2n(\mathrm{d}e) \int_0^{\sigma(e)} \frac{\mathrm{d}a}{(x + e_a)^2}\right),$$

where $n(\mathrm{d}e)$ is the Itô measure of positive excursions of Brownian motion (so that $2n$ is the excursion measure of the reflected Brownian motion $\hat{\beta}$), and $\sigma(e)$ is the lifetime of the generic excursion e . The Bismut decomposition of the measure n finally gives

$$\int n(\mathrm{d}e) \int_0^{\sigma(e)} \frac{\mathrm{d}a}{(x + e_a)^2} = \int_0^\infty \frac{\mathrm{d}r}{(x + r)^2} = \frac{1}{x},$$

hence (18). Putting (16), (17) and (18) together, we deduce that

$$\begin{aligned} \left(1 - \mathbb{P}(\Delta'_0 > xm\sqrt{\frac{2}{3}})\right)^{m+1} &= \mathbb{P}\left(\frac{1}{m}\sqrt{\frac{3}{2}} \max\{\Delta'_i : 0 \leq i \leq m\} \leq x\right) \\ &= \mathbb{P}\left(\frac{1}{m}\sqrt{\frac{3}{2}} \max\{-V'(j) + \underline{X}_{-\underline{C}(j)} : 0 \leq i \leq \kappa'(m)\} \leq x\right) \\ &\xrightarrow{m \rightarrow \infty} \mathbb{P}\left(\sup\{Z_t - \hat{\beta}_{-\underline{B}_t} : 0 \leq t \leq T'_{\sqrt{3/2}}\} \leq x\right) = \exp(-\sqrt{6}/x), \end{aligned}$$

where at the penultimate step we used the fact that the law of $\sup\{Z_t - \hat{\beta}_{-\underline{B}_t} : 0 \leq t \leq T'_{\sqrt{3/2}}\}$ is diffuse, and the fact that $(B, -Z, -\beta)$ and (B, Z, β) have the same distribution. Therefore, taking $x = \sqrt{3/2}$, we get

$$\mathbb{P}(\Delta'_0 > m) \sim \frac{2}{m},$$

concluding the proof of Lemma 14. \square

5 Horoballs and points of escape to infinity

In the two remaining sections, we use the representation of the UIPQ given by Theorem 1 in order to deduce new results on this object. First, inspired by the work of Krikun [20], we study the length of the separating cycle around the origin of the UIPQ at “height” $-r$ seen from infinity. In the following, we let Q_∞ be a random variable with the law of the UIPQ, and we let $((T_\infty, \ell), \eta)$ be the labeled tree associated with Q_∞ by the Schaeffer correspondence.

For every integer $l \in \mathbb{Z}$, we denote by $H_l = H_l(Q_\infty) = \{v \in V(Q_\infty) : \ell(v) \leq l\}$. In view of Theorem 1, we can interpret H_l as the *ball centered at infinity*, or *horoball*, with “radius $\mathrm{d}_{\mathrm{gr}}^{Q_\infty}(\emptyset, \infty) + l$ ”, where ∞ is a point at infinity in Q_∞ . Intuitively, the boundary of this set (a “horosphere”) is made of several disjoint cycles, one of which separates \emptyset from ∞ . We are going to give asymptotic properties for the length of this cycle, the set of “points to escape to ∞ at level l ”.

To this purpose, it is easier to work with a slightly modified graph \hat{Q}_∞ , which is obtained by adding to Q_∞ the edges of T_∞ given by the inverse Schaeffer construction of Section 2.3.2. Therefore, in faces f of Q_∞ , around which the four vertices in clockwise order have labels $l, l+1, l, l+1$, we add the diagonal between the vertices with label $l+1$ (such faces are called confluent faces in [12]). In faces with labels $l, l+1, l+2, l+1$, we just double the edge between the last two vertices. The map \hat{Q}_∞ is no longer a quadrangulation: Some of its faces are still squares, but others are triangles, and some others have degree 2. Nonetheless, Q_∞ and \hat{Q}_∞ are very similar: They have the same vertex set, and it is easy to check that formula (7) remains true in this context, i.e.

$$\ell(v) = \lim_{z \rightarrow \infty} \left(\mathrm{d}_{\mathrm{gr}}^{Q_\infty}(z, \emptyset) - \mathrm{d}_{\mathrm{gr}}^{Q_\infty}(z, v) \right) = \lim_{z \rightarrow \infty} \left(\mathrm{d}_{\mathrm{gr}}^{\hat{Q}_\infty}(z, \emptyset) - \mathrm{d}_{\mathrm{gr}}^{\hat{Q}_\infty}(z, v) \right)$$

for every vertex v , because we only added edges between vertices with the same label, and geodesic rays in \hat{Q}_∞ never use the new edges. We are not going to need this fact in the sequel, so details are left to the reader.

Now the complement $\{v \in V(Q_\infty) : \ell(v) > l\}$ of H_l induces a subgraph of \hat{Q}_∞ , and if $l < 0$ we let $F_l = F_l(Q_\infty)$ be the set of vertices in the connected component of this subgraph that contains \emptyset . We let ∂F_l be the set of vertices of H_l that are connected to F_l by an edge. Yet otherwise said, F_l is the set of vertices v with $\ell(v) > l$, and which can be joined to \emptyset by a path of \hat{Q}_∞ along which labels are all strictly greater than l , and ∂F_l is the set of vertices with label l that can be joined to \emptyset by a path in Q_∞ along which all labels are all strictly greater than l except at the initial point.

Although it is not obvious at first sight, the set F_l is almost-surely finite: The next statement implies that it is contained in the set of vertices of the subtrees $L_0, \dots, L_{\sigma_l}, R_0, \dots, R_{\sigma_l}$ to the left and to the right of T_∞ , where $\sigma_l = \inf\{n \geq 0 : \ell(S(n)) = -l\}$ is a.s. finite. Recall that $[[a, b]]$ is the path from a to b in T_∞ .

Proposition 18. *Let $r > 0$ and $v \in V(Q_\infty)$. Then*

1. *v belongs to $F_{-r}(Q_\infty)$ if and only if $\ell(v') > -r$ for every $v' \in [[\emptyset, v]]$, and*
2. *v belongs to $\partial F_{-r}(Q_\infty)$ if and only if $\ell(v) = -r$ and $\ell(v') > -r$ for every $v' \in [[\emptyset, v]] \setminus \{v\}$.*

Proof. Let v be such that $\ell(v') > -r$ for every $v' \in [[\emptyset, v]] \setminus \{v\}$. The path $[[\emptyset, v]]$ in T_∞ is also a path in the augmented graph \hat{Q}_∞ , and it goes only through vertices outside H_{-r} , except maybe at v . So if furthermore $\ell(v) = -r$ we obtain that $v \in \partial F_{-r}$, otherwise $v \in F_{-r}$.

Conversely, suppose that there is a vertex v' on the path $[[\emptyset, v]] \setminus \{v\}$ with $\ell(v') \leq -r$. We consider the two corners $c_{(1)}$ and $c_{(2)}$ that are respectively the smallest and largest corner incident to the vertex v' . We then construct the geodesics γ_1 and γ_2 emanating from the corners $c_{(1)}$ and $c_{(2)}$ by taking consecutive successors $\mathcal{S}^{(i)}(c_{(1)}), i \geq 0$ and $\mathcal{S}^{(i)}(c_{(2)}), i \geq 0$ respectively.

The geodesics γ_1 and γ_2 coalesce at the first corner c following $c_{(2)}$ in the contour with label $\ell_{\min} - 1$, where ℓ_{\min} is the smallest label of the corners between $c_{(1)}$ and $c_{(2)}$. The concatenation of the parts of γ_1 and γ_2 between v' and $\mathcal{V}(c)$ induces a cycle \mathcal{C} of the map Q_∞ , such that $v \notin \mathcal{C}$ and separates v from \emptyset in Q_∞ , see Figure 11 for an example. Note that all the vertices of \mathcal{C} have labels less than or equal to $-r$. By the Jordan Theorem, any path in \hat{Q}_∞ joining \emptyset to v crosses \mathcal{C} , and thus has a vertex, other than v , with label less than or equal to $-r$. It follows that v does not belong to $\partial F_{-r} \cup F_{-r}$. \square

We can use Proposition 18 to derive asymptotics for the number of vertices in $\partial F_{-r}(q)$:

Theorem 3. *The sequence $(2|\partial F_{-r}(Q_\infty)|/r^2, r \geq 1)$ converges in distribution to a random variable W with Laplace transform*

$$E[e^{-\lambda W}] = \left(\frac{1}{1 + \sqrt{\lambda}} \right)^3.$$

Remark. A similar problem has already been studied by Krikun. In [20], he considered the component $\partial \tilde{F}_r(Q_\infty)$ of the boundary of the ball $B_{\mathbf{Q}, r}(Q_\infty)$ that separates the root vertex from infinity. He showed that $2|\partial \tilde{F}_r(Q_\infty)|/r^2$ converges in distribution as $r \rightarrow \infty$ to a standard Gamma random variable with parameter $3/2$. In our setup, $|\partial F_{-r}(Q_\infty)|$ is, roughly speaking, the length of the boundary of the horoball that separates the root from infinity at level $-r$. It is not surprising that the typical size of this component should be of order r^2 as well, and obey a similar limiting result. However, we see that the limiting distribution differs from the Gamma law. In this respect, it is interesting to compare the tail distributions of these variables. Let W, \tilde{W} have respective Laplace transforms

$$\phi(\lambda) = E[e^{-\lambda W}] = \left(\frac{1}{1 + \sqrt{\lambda}} \right)^3, \quad \tilde{\phi}(\lambda) = E[e^{-\lambda \tilde{W}}] = \left(\frac{1}{1 + \lambda} \right)^{3/2},$$




Figure 11: Cycles separating vertices from \emptyset composed by two geodesics.

so that \tilde{W} follows the Gamma distribution with parameter $3/2$. As $\lambda \rightarrow \infty$, we have $\phi(\lambda) \sim \tilde{\phi}(\lambda) \sim \lambda^{-3/2}$, so that a Tauberian Theorem [8, Theorem 1.7.1] entails that as $y \rightarrow 0^+$

$$\mathbb{P}(W \leq y) \sim \mathbb{P}(\tilde{W} \leq y) \sim \frac{4}{3\sqrt{\pi}} y^{3/2},$$

so these two distributions have a similar behavior at 0. By contrast, \tilde{W} has exponential tails, while we have

$$\phi(\lambda) = 1 - 3\sqrt{\lambda} + o(\sqrt{\lambda}), \quad \text{when } \lambda \rightarrow 0$$

which by applying Corollary 8.1.7 in [8] shows that

$$\mathbb{P}(W \geq y) \sim \frac{3}{\sqrt{\pi}} y^{-1/2},$$

as $y \rightarrow \infty$. Therefore, W has a heavy tail.

Proof of Theorem 3. Assume that Q_∞ is obtained by the extended Schaeffer bijection, that is $Q_\infty = \Phi((T_\infty, \ell), \eta)$ where $((T_\infty, \ell), \eta)$ has law $\mu \otimes \mathcal{B}(1/2)$. The only vertex of $\partial F_{-r}(Q_\infty)$ that belongs to the spine of T_∞ is $S(\sigma_r)$ where we recall that $\sigma_r = \inf\{i \geq 0 : \ell(S(i)) = -r\}$. Then, Proposition 18 implies that

$$|\partial F_{-r}(Q_\infty)| = 1 + \sum_{i < \sigma_r} |Y_{L_i}(-r)| + \sum_{i < \sigma_r} |Y_{R_i}(-r)| \quad (19)$$

where, if θ is a labeled tree whose root label is strictly larger than $-r$, $Y_\theta(-r)$ is the set of all vertices of θ with label $-r$ and such that all their ancestors have a label strictly larger than $-r$. Recalling that the trees (L_i) and (R_i) are independent conditionally given (X_i) , we get from

(19) that

$$\begin{aligned}
& E \left[\exp -\lambda \frac{2|\partial F_{-r}(Q\infty)|}{r^2} \right] \\
&= \exp \left(-\frac{2\lambda}{r^2} \right) E \left[\prod_{i < \sigma_r} E \left[\exp \left(-\frac{2\lambda}{r^2} |Y_{L_i}(-r)| \right) \middle| X_i \right] E \left[\exp \left(-\frac{2\lambda}{r^2} |Y_{R_i}(-r)| \right) \middle| X_i \right] \right] \\
&= \exp \left(-\frac{2\lambda}{r^2} \right) E \left[\prod_{i < \sigma_r} \rho_{X_i} \left(\exp \left(-\frac{2\lambda}{r^2} |Y_\theta(-r)| \right) \right)^2 \right] \\
&= \exp \left(-\frac{2\lambda}{r^2} \right) E \left[\exp \left(2 \sum_{i < \sigma_r} \log \rho_{X_i} \left(\exp \left(-\frac{2\lambda}{r^2} |Y_\theta(-r)| \right) \right) \right) \right]. \tag{20}
\end{aligned}$$

To compute the right hand side of (20), we need to evaluate the generating functions:

$$f_{l,-r}(x) = \rho_l \left(x^{|Y_\theta(-r)|} \right), \quad 0 \leq x \leq 1$$

for $l \geq -r$, with the boundary condition $f_{-r,-r}(x) = x$. The measures ρ_l being the laws of Galton Watson trees with geometric(1/2) offspring distribution and uniform labels, it is easy to derive the following recursive relation for $l > -r$:

$$\begin{aligned}
f_{l,-r}(x) &= \frac{1}{2} \sum_{k=0}^{\infty} \sum_{i_1+i_2+i_3=k} \binom{k}{i_1, i_2, i_3} \left(\frac{1}{6} f_{l-1,-r}(x) \right)^{i_1} \left(\frac{1}{6} f_{l,-r}(x) \right)^{i_2} \left(\frac{1}{6} f_{l+1,-r}(x) \right)^{i_3} \\
&= \frac{1}{2} \left(1 - \frac{f_{l-1,-r}(x) + f_{l,-r}(x) + f_{l+1,-r}(x)}{6} \right)^{-1}.
\end{aligned}$$

From this identity, we get the following recurrence relation for $l > -r$:

$$2f_{l,-r}(x) = 1 + \frac{1}{12} 2f_{l,-r}(x) (2f_{l-1,-r}(x) + 2f_{l,-r}(x) + 2f_{l+1,-r}(x)).$$

To solve this equation we follow [9]. Putting

$$F(x, y) = xy \left(1 - \frac{1}{12}x - \frac{1}{12}y \right) - x - y,$$

then $F(2f_{l,-r}(x), 2f_{l+1,-r}(x))$ does not depend on $x \in [0, 1]$ and $l \geq -r$, since

$$\begin{aligned}
& F(2f_{l,-r}(x), 2f_{l+1,-r}(x)) - F(2f_{l-1,-r}(x), 2f_{l,-r}(x)) = (2f_{l+1,-r}(x) - 2f_{l-1,-r}(x)) \\
& \quad \times \left(2f_{l,-r}(x) - 1 - \frac{1}{12} 2f_{l,-r}(x) (2f_{l-1,-r}(x) + 2f_{l,-r}(x) + 2f_{l+1,-r}(x)) \right).
\end{aligned}$$

It is easy to verify that $f_{l,-r}(x) \rightarrow 1$ as $l \rightarrow \infty$ and since $F(2, 2) = -\frac{4}{3}$, we have the relation:

$$F(2f_{l,-r}(x), 2f_{l+1,-r}(x)) = -\frac{4}{3}, \tag{21}$$

for $l \geq -r$, with the initial condition $f_{-r,-r}(x) = x$. The general solution of (21) is given by

$$f_{l,-r} = 1 - \frac{2}{(l+r+a)(l+r+1+a)}, \tag{22}$$

for $l \geq -r$, where $a = a(x)$ is a function, which from the initial condition is found to be

$$a(x) = \frac{-1 + \sqrt{1 + 8(1-x)^{-1}}}{2},$$

for $x \in [0, 1]$.

Substituting (22) in (20), one gets:

$$\begin{aligned}
& \exp\left(\frac{2\lambda}{r^2}\right) E\left[\exp -2\lambda \frac{|\partial F_{-r}(Q_\infty)|}{r^2}\right] \\
&= E\left[\exp 2 \sum_{i=0}^{\sigma_r-1} \log\left(1 - \frac{2}{(X_i + r + a(e^{-2\lambda/r^2})) (X_i + r + 1 + a(e^{-2\lambda/r^2}))}\right)\right] \\
&= E\left[\exp 2r^2 \int_0^{\sigma_r/r^2} dt \log\left(1 - \frac{2}{(X_{\lfloor r^2 t \rfloor} + r + a(e^{-2\lambda/r^2})) (X_{\lfloor r^2 t \rfloor} + r + 1 + a(e^{-2\lambda/r^2}))}\right)\right]. \tag{23}
\end{aligned}$$

By Skorokhod's representation theorem we can find a sequence of processes $((X_k^{(r)})_{k \geq 0}, r \geq 0)$ such that for each $r \geq 1$ we have $(X_k^{(r)})_{k \geq 0} = (X_k)_{k \geq 0}$ in law and such that we have the following almost sure convergence

$$\left(\frac{1}{r} \sqrt{\frac{3}{2}} X_{\lfloor r^2 t \rfloor}^{(r)}\right)_{t \geq 0} \xrightarrow[r \rightarrow \infty]{a.s.} (\beta_t)_{t \geq 0},$$

where β is a standard Brownian motion. It is also easy to check that, as $r \rightarrow \infty$, one has $r^{-1}a(e^{-2\lambda/r^2}) \rightarrow 1/\sqrt{\lambda}$. This gives

$$\begin{aligned}
& r^2 \log\left(1 - \frac{2}{(X_{\lfloor r^2 t \rfloor}^{(r)} + r + a(e^{-2\lambda/r^2})) (X_{\lfloor r^2 t \rfloor}^{(r)} + r + 1 + a(e^{-2\lambda/r^2}))}\right) \\
&= r^2 \log\left(1 - \frac{1}{r^2} \frac{2}{(X_{\lfloor r^2 t \rfloor}^{(r)}/r + 1 + a(e^{-2\lambda/r^2})/r)^2} + o\left(\frac{1}{r^2}\right)\right) \\
&\longrightarrow -\frac{2}{\left(\sqrt{\frac{2}{3}}\beta_t + 1 + 1/\sqrt{\lambda}\right)^2}
\end{aligned}$$

almost surely as $r \rightarrow \infty$. Furthermore, if we denote $\sigma_r^{(r)}$ the first hitting time of $-r$ by the process $X^{(r)}$, then one has the almost sure convergence:

$$\frac{\sigma_r^{(r)}}{r^2} \xrightarrow[r \rightarrow \infty]{} T_{\sqrt{3/2}}(B),$$

where $T_{\sqrt{3/2}}(B)$ is the first hitting time of $-\sqrt{3/2}$ of the Brownian motion B . This is an easy consequence of the fact that almost-surely, B takes values strictly less than $-3/2$ on any time-interval of the form $[T_{\sqrt{3/2}}(B), T_{\sqrt{3/2}}(B) + \varepsilon]$ for $\varepsilon > 0$. Thus replacing X by $X^{(r)}$ into (23), an argument of dominated convergence then gives

$$E\left[\exp -\lambda \frac{2|\partial F_{-r}(Q_\infty)|}{r^2}\right] \xrightarrow[r \rightarrow \infty]{} E\left[\exp\left(-4 \int_0^{T_{\sqrt{3/2}}(B)} \frac{du}{\left(\sqrt{\frac{2}{3}}\beta_u + 1 + \sqrt{1/\lambda}\right)^2}\right)\right],$$

and the scaling property of the Brownian motion shows that the right hand side of the last display is equal to

$$E\left[\exp -6 \int_0^{T_1(B)} \frac{dt}{(B_t + 1 + \sqrt{1/\lambda})^2}\right].$$

Let us write \mathbb{P}_x for the law of $(x + B_t, t \geq 0)$, and let $(Z_t, t \geq 0)$ be the canonical process. Let also $T_y(Z) = \inf\{t \geq 0 : Z_t = y\} \in [0, \infty]$ be the first hitting time of y . By translation, we can re-write the previous expectation as

$$\mathbb{E}_{1+\sqrt{1/\lambda}} \left[\exp \left(-6 \int_0^{T_{\sqrt{1/\lambda}}(Z)} \frac{dt}{Z_t^2} \right) \right].$$

At this point, we can use the absolute continuity relations between Bessel processes with different indices, due to Yor [33, Exercise XI.1.22] (see also [28] for a similar use of these absolute continuity relations). The last expectation then equals

$$\left(\frac{1 + \sqrt{1/\lambda}}{\sqrt{1/\lambda}} \right)^4 \mathbb{P}_{1+\sqrt{1/\lambda}}^{(9)}(T_{\sqrt{1/\lambda}} < \infty),$$

where $\mathbb{P}_x^{(9)}$ is the law of the 9-dimensional Bessel process started from $x > 0$. It is classical that $\mathbb{P}_x^{(9)}(T_y < \infty) = (y/x)^7$ for every positive x, y with $y < x$. This can be verified from the fact that $(Z_t^{-7}, t \geq 0)$ is a local martingale under $\mathbb{P}_x^{(9)}$, as can be checked from Ito's formula, and the fact that Z under $\mathbb{P}_x^{(9)}$ has same distribution as the Euclidean norm of a 9-dimensional Brownian motion started from a point with norm x . We finally obtain that

$$E \left[\exp -6 \int_0^{T_1(B)} \frac{dt}{(\beta_t + 1 + \sqrt{1/\lambda})^2} \right] = \left(\frac{1}{1 + \sqrt{\lambda}} \right)^3,$$

as wanted. □

6 Random walk on the UIPQ

This section focuses on the simple random walk over the UIPQ. We first provide a proof of a known fact (see [21]) that the distribution of the UIPQ is invariant under re-rooting along a simple random walk. We then make a step in understanding the recurrence/transience property of the walk on the UIPQ.

6.1 Invariance under re-rooting along the random walk

Let q be a rooted quadrangulation, which can be finite or infinite. We consider the nearest-neighbor random walk on q starting from e_+^* . Rather than the random sequence of vertices visited by this walk, we really want to emphasize the sequence of edges that are visited. Formally, we consider a random infinite sequence of oriented edges (E_0, E_1, E_2, \dots) starting with the root edge $E_0 = e^*$ and defined recursively as follows. Conditionally given $(E_i, 0 \leq i \leq j)$, we let E_{j+1} be a random edge pointing from $(E_j)_+$, chosen uniformly among the $\deg((E_j)_+)$ possible ones. The sequence $((E_{1+i})_-, i \geq 0)$ is then the usual nearest-neighbor random walk on $V(q)$, starting from e_+^* .

We let P_q be the law of the sequence $(E_i, i \geq 0)$ ¹. Also, for any oriented edge e of the map q , we let $q^{(e)}$ be the map q re-rooted at e . Finally, if λ is a probability distribution on \mathbf{Q} , let $\Theta^{(r)}(\lambda)$ be the probability distribution defined by

$$\Theta^{(r)}(\lambda)(A) = \int_{\mathbf{Q}} \lambda(dq) \int P_q(d(e_0, e_1, e_2, \dots)) \mathbf{1}_{q^{(e_r)} \in A},$$

¹Recall that a map is an equivalence class of embedded graphs, so the last definition does not really make sense but the reader can check that all quantities computed in the sequel do not depend on a representative embedded graph of the map.

for any Borel subset A of \mathbf{Q} . The probability measure $\Theta^{(r)}(\lambda)$ is the distribution of a random map with distribution λ , re-rooted at the r th step of the random walk.

Proposition 19. *The law ν of the UIPQ is invariant under re-rooting along a simple random walk, in the sense that for every $r \geq 0$, one has $\Theta^{(r)}(\nu) = \nu$.*

Moreover, if A is an event of the Borel σ -algebra of $(\mathbf{Q}, d_{\mathbf{Q}})$ such that $\nu(A) = 1$, then

$$\nu \left(\left\{ q \in \mathbf{Q} : \forall e \in \vec{E}(q), q^{(e)} \in A \right\} \right) = 1.$$

See [1, 6] for a general study of random graphs that are invariant under re-rooting along the simple random walk. In the case of the UIPQ, the first assertion of Proposition 19 appears in [21, Section 1.3], see also [4, Theorem 3.2] for a similar result in the case of the UIPT. We provide a detailed proof for the sake of completeness.

Proof. It is easy to see that the function $\Theta^{(r)}$ on the set $\mathcal{P}(\mathbf{Q})$ of Borel probability measures on $(\mathbf{Q}, d_{\mathbf{Q}})$ coincides with the r -fold composition of $\Theta = \Theta^{(1)}$ with itself. Therefore, it suffices to show the result for $r = 1$.

Let us check that Θ is continuous when $\mathcal{P}(\mathbf{Q})$ is endowed with the topology of weak convergence. Indeed, if λ_n converges weakly to λ as $n \rightarrow \infty$, then by the Skorokhod representation theorem, we can find a sequence $(Q_n, n \geq 0)$ of random variables in \mathbf{Q} with respective laws $(\lambda_n, n \geq 0)$, that converges a.s. to a random variable Q with law λ . For every fixed $R > 0$, it then holds that $B_{\mathbf{Q}, R}(Q_n) = B_{\mathbf{Q}, R}(Q)$ for every n large enough a.s.. Now, we can couple in an obvious way the random walks with laws P_{Q_n} and P_Q , in such a way that the first step E_1 is the same edge in Q_n and Q on the event where $B_{\mathbf{Q}, 1}(Q_n) = B_{\mathbf{Q}, 1}(Q)$. For such a coupling, we then obtain that $B_{\mathbf{Q}, R-1}(Q_n^{(E_1)}) = B_{\mathbf{Q}, R-1}(Q^{(E_1)})$ for every n large enough. Since R is arbitrary, this shows that $Q_n^{(E_1)}$ converges a.s. to $Q^{(E_1)}$, so that $\Theta(\lambda_n)$ converges weakly to $\Theta(\lambda)$, as desired.

Since we know by Theorem 2.1.2 that the uniform law ν_n on \mathbf{Q}_n converges to ν , it suffices to show that $\Theta(\nu_n) = \nu_n$. Now consider the law of the doubly-rooted map (q, e^*, e_1) under the law $\nu_n(dq)P_q(d(e_i)_{i \geq 0})$. The probability that (q, e^*, e_1) equals a particular doubly-rooted map (q, e', e'') with $e'_+ = e''_-$ is equal to $(\#\mathbf{Q}_n \deg(e'_+))^{-1}$, from which it immediately follows that (q, e_*, e_1) has the same distribution as $(q, \overleftarrow{e}_1, \overleftarrow{e}_*)$, still under $\nu_n(dq)P_q(d(e_i)_{i \geq 0})$. Hence (q, \overleftarrow{e}_1) under $\nu_n(dq)P_q(d(e_i)_{i \geq 0})$ has the same law ν_n as (q, e_*) . Since ν_n is obviously invariant under the reversal of the root edge, we get that (q, e_1) has law ν_n . But by definition, it also has law $\Theta(\nu_n)$, which gives the first assertion of Proposition 19.

Let us now prove the last part of the statement of the proposition. By the first part, we have

$$\int_{\mathbf{Q}} \nu(dq) E_q \left[\sum_{n=0}^{\infty} \mathbf{1}_{A^c}(q^{(e_n)}) \right] = 0.$$

Thus, $\nu(dq)$ a.s. , $E_q[\sum_{n=0}^{\infty} \mathbf{1}_{A^c}(q^{(e_n)})] = 0$. But

$$E_q \left[\sum_{n=0}^{\infty} \mathbf{1}_{A^c}(q^{(e_n)}) \right] \geq \sum_{e \in \vec{E}(q)} P_q(\exists n \geq 0 : e_n = e) \mathbf{1}_{A^c}(q^{(e)}),$$

and $P_q(\exists n \geq 0 : e_n = e) > 0$ for every $e \in \vec{E}(q)$ because q is connected. This completes the proof. \square

Remark. It can seem a little unnatural to fix the first step of the random walk to be equal to e^* , hence to be determined by the rooted map q rather than by some external source of randomness. In fact, we could also first re-root the map at some uniformly chosen random edge incident to e^* , and start the random walk with this new edge. Since the first re-rooting leaves the laws ν_n, ν invariant, as is easily checked along the same lines as the previous proof, the results of Proposition 19 still hold with the new random walk.

6.2 On recurrence

Let Q_∞ be the uniform infinite planar quadrangulation. Conditionally on Q_∞ , $(E_k)_{k \geq 0}$ denotes the random sequence of oriented edges with $E_0 = e^*$ traversed by a simple random walk on Q_∞ as discussed at the beginning of Section 6.1. We write $\mathcal{X}_k = (E_k)_-$ for the sequence of vertices visited along the walk. For $k \geq 0$, we denote the quadrangulation Q_∞ re-rooted at the oriented edge E_k by $Q_\infty^{(k)}$. Proposition 19 shows that $Q_\infty^{(k)}$ has the same distribution as Q_∞ .

Question 20 ([4]). *Is the simple random walk $(\mathcal{X}_k)_{k \geq 0}$ on Q_∞ almost surely recurrent?*

A similar question for UIPT arose when Angel & Schramm [4] introduced this infinite random graph. These questions are still open. James T. Gill and Steffen Rohde [17] proved that the Riemann surface obtained from the UIPQ by gluing squares along edges is recurrent for Brownian motion. The first author and Itai Benjamini also proved that the UIPQ is almost surely Liouville [6]. However the lack of a bounded degree property for the UIPQ prevents one from deducing recurrence from these results (see also [7]). Our new construction of the UIPQ however leads to some new information suggesting that the answer to the above Question should be positive.

Theorem 4. *The process $(\ell(\mathcal{X}_n))_{n \geq 0}$ is a.s. recurrent, i.e. visits every integer infinitely often.*

Proof. For every $k \geq 0$, one can consider the labeling $(\ell^{(k)}(u))_{u \in Q_\infty}$ of the vertices of Q_∞ that corresponds to the labeling given by Theorem 1 applied to the rooted infinite planar quadrangulation $Q_\infty^{(k)}$. On the one hand, it is straightforward to see from (6) that $\ell^{(k)}(u) - \ell^{(k)}(v) = \ell(u) - \ell(v)$ for every $u, v \in Q_\infty$. On the other hand, applying Proposition 19 we deduce that the process $(\ell^{(k)}(\mathcal{X}_{k+i}) - \ell^{(k)}(\mathcal{X}_k))_{i \geq 0}$ has the same distribution as $(\ell(\mathcal{X}_i) - \ell(\mathcal{X}_0))_{i \geq 0}$. Gathering up the pieces, we deduce that for every integer $k \geq 0$ we have

$$(\ell(\mathcal{X}_i) - \ell(\mathcal{X}_0))_{i \geq 0} \stackrel{(d)}{=} (\ell(\mathcal{X}_{k+i}) - \ell(\mathcal{X}_k))_{i \geq 0}. \quad (24)$$

Hence the increments $(\ell(\mathcal{X}_{i+1}) - \ell(\mathcal{X}_i))_{i \geq 0}$ form a stationary sequence. Furthermore, we have $|\ell(\mathcal{X}_1) - \ell(\mathcal{X}_0)| = 1$, and since the distribution of Q_∞ is preserved when reversing the orientation of the root edge we deduce

$$\ell(\mathcal{X}_1) - \ell(\mathcal{X}_0) \stackrel{(d)}{=} \ell(\mathcal{X}_0) - \ell(\mathcal{X}_1) \stackrel{(d)}{=} \mathcal{B}(1/2).$$

In particular the increments of $\ell(\mathcal{X}_n)$ have zero mean. Suppose for an instant that the increments of $\ell(\mathcal{X}_n)$ were also ergodic, then Theorem 3 of [15] would directly apply and give the recurrence of $\ell(\mathcal{X}_n)$. Although the UIPQ is ergodic, a proof of this fact would take us too far, so we will reduce the problem to the study of ergodic components.

By standard facts of ergodic theory, the law ξ of the sequence of increments $(\ell(\mathcal{X}_{i+1}) - \ell(\mathcal{X}_i))_{i \geq 0}$ can be expressed as a barycenter of ergodic probability measures in the sense of Choquet, namely for every $A \subset \mathcal{B}(\mathbb{R})^{\otimes \mathbb{N}}$ we have

$$\xi(A) = \int \zeta(A) dm(\zeta), \quad (25)$$

where m is a probability measure on the set of all probability measures on $(\mathbb{R}^{\mathbb{N}}, \mathcal{B}(\mathbb{R})^{\otimes \mathbb{N}}, \mathbb{P})$ that are ergodic for the shift. In our case, it suffices to show that m -almost every ζ satisfies the assumption of [15, Theorem 3]. Specializing (25) with $A_1 = \{(y_i)_{i \geq 0} : |y_{i+1} - y_i| \leq 1, \forall i \geq 0\}$ we deduce that m -almost every ζ , we have $\zeta(A_1) = 1$, in particular the increments under ζ are integrable. It remains to show that they have zero mean.

Lemma 21. *Almost surely we have*

$$\lim_{n \rightarrow \infty} \frac{\ell(\mathcal{X}_n)}{n} = 0.$$

Proof. In [11, Theorem 6.4] it is shown that $\mathbb{E}[\#B_{\mathbf{Q},r}(Q_\infty)] \leq C_3 r^4$ where $C_3 > 0$ is independent of $r \geq 1$. Using the Borel-Cantelli lemma we easily deduce that

$$\lim_{r \rightarrow \infty} r^{-6} \#B_{\mathbf{Q},r}(Q_\infty) = 0, \quad a.s. \quad (26)$$

We now use the classical Varopoulos-Carne upper bound (see for instance Theorem 13.4 in [29]): we have

$$p_n(e_+^*, x) \leq 2 \sqrt{\frac{\deg(x)}{\deg(e_+^*)}} \exp\left(-\frac{d_{\text{gr}}^{Q_\infty}(e_+^*, x)^2}{2n}\right), \quad (27)$$

where conditionally on Q_∞ , $p_n(\cdot, \cdot)$ is the n -step transition probability of the simple random walk started from e_+^* in Q_∞ . Conditionally on Q_∞ , using a crude bound $\deg(x) \leq \#B_{\mathbf{Q},n+1}(Q_\infty)$ on the degree of a vertex $x \in B_{\mathbf{Q},n}(Q_\infty)$, we have using (27)

$$P_{Q_\infty}(\mathcal{X}_n \notin B_{\mathbf{Q},n^{2/3}}(Q_\infty)) \leq 2 \exp\left(-\frac{n^{1/3}}{2}\right) (\#B_{\mathbf{Q},n+1}(Q_\infty))^{3/2}.$$

Hence on the event $\{\lim_{r \rightarrow \infty} r^{-6} \#B_{\mathbf{Q},r}(Q_\infty) = 0\}$, an easy application of the Borel-Cantelli lemma shows that $n^{-1} d_{\text{gr}}^{Q_\infty}(\mathcal{X}_n, \emptyset) \rightarrow 0$ as $n \rightarrow \infty$. Since $|\ell(\mathcal{X}_n)| \leq d_{\text{gr}}^{Q_\infty}(\mathcal{X}_n, \emptyset)$, the above discussion together with (26) completes the proof of the lemma. \square

Let us complete the proof of Theorem 4. We can specialize formula (25) to $A_2 = \{(y_i)_{i \geq 0} : \lim i^{-1}|y_i| = 0\}$, to obtain that m -a.e ζ we have $\zeta(A_2) = 1$. Using the ergodic theorem that means that the increments under ζ are centered. We can thus apply Theorem 3 of [15] to get that for m -almost every ζ , the process whose increments are distributed according to ζ is recurrent, hence $(\ell(\mathcal{X}_n))$ is almost surely recurrent. \square

Appendix: infinite maps and their embeddings

In this section, we explain how the elements of \mathbf{Q}_∞ can be seen as infinite quadrangulations of a certain non-compact surface, completing the description of Section 2.1.1.

Recall that an element q of \mathbf{Q}_∞ is a sequence of compatible maps with holes (q_1, q_2, \dots) , in the sense that $q_r = B_{\mathbf{Q},r}(q_{r+1})$. This sequence defines a unique cell complex S_q up to homeomorphism, with an infinite number of 2-cells, which are quadrangles. This cell complex is an orientable, connected, separable topological surface, and every compact connected sub-surface is planar.

It is known [34] that the topology of S_q is characterized by its *ends space*, which is a certain totally disconnected compact space. Roughly speaking, the ends space determines the different “points at infinity” of the surface. More precisely, following [34], we define a boundary component of S_q as a sequence (U_1, U_2, \dots) of subsets of S_q , such that

- for every $i \geq 1$, the set U_i is unbounded, open, connected and with compact boundary,
- for every $i \geq 1$, it holds that $U_{i+1} \subset U_i$,
- for every bounded subset $A \subset S_q$, $U_i \cap A = \emptyset$ for every i large enough.

Two boundary components $(U_i, i \geq 1), (U'_i, i \geq 1)$ are called equivalent if for every $i \geq 1$ there exists $i' \geq 1$ such that $U'_{i'} \subset U_i$, and vice-versa. An *end* is an equivalence class of boundary components. For every $U \subset S_q$ with compact boundary, we let V_U be the set of all ends whose corresponding boundary components are sequences of sets which are eventually included in U . The topological space having the sets V_U as a basis is called the ends space, and denoted by \mathcal{E}_q .

Conversely, it is plain that every rooted quadrangulation of an orientable, connected, separable, non-compact planar surface, defines an element of \mathbf{Q}_∞ , by taking the sequence of the balls centered at the root vertex, with the same definition as in Section 2.1.1. The separability ensures that the collection of balls exhausts the whole surface. Thus we have:

Proposition 22. *The elements of \mathbf{Q}_∞ are exactly the quadrangulations of orientable, connected, separable, non-compact planar surfaces, and considered up to homeomorphisms that preserve the orientation.*

To understand better what the ends space is in our context, note that there is a natural tree structure \mathcal{T}_q associated with $q \in \mathbf{Q}_\infty$. The vertices v of this tree are the holes of q_1, q_2, q_3, \dots , and an edge links the vertices v and v' if there exists an $r \geq 1$ such that v is a hole of q_r , v' is a hole of q_{r+1} , and v' is included in the face determined by v . Furthermore, all the holes in q_1 are linked by an edge to an extra root vertex.

It is then easy to see that \mathcal{E}_q is homeomorphic to the ends space $\partial\mathcal{T}_q$ which is defined as follows: $\partial\mathcal{T}_q$ is just the set of infinite injective paths (spines) in \mathcal{T}_q starting from the root, and a basis for its topology is given by the sets W_v made of the spines that pass through the vertex v of \mathcal{T}_q . (This is consistent, since it is easy and well-known that the ends space of trees with finite degrees is a compact totally disconnected space.)

In particular, when \mathcal{T}_q has a unique spine, then \mathcal{E}_q is reduced to a point, which means that the topology of S_q is that of the plane \mathbb{R}^2 .

References

- [1] D. Aldous and R. Lyons. Processes on unimodular random networks. *Electron. J. Probab.*, 12:no. 54, 1454–1508 (electronic), 2007.
- [2] J. Ambjørn, B. Durhuus, and T. Jonsson. *Quantum geometry*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1997. A statistical field theory approach.
- [3] O. Angel. Growth and percolation on the uniform infinite planar triangulation. *Geom. Funct. Anal.*, 13(5):935–974, 2003.
- [4] O. Angel and O. Schramm. Uniform infinite planar triangulation. *Comm. Math. Phys.*, 241(2-3):191–213, 2003.
- [5] I. Benjamini. Random planar metrics. *Proceedings of the ICM 2010*, 2010.
- [6] I. Benjamini and N. Curien. Ergodic theory on stationary random graphs. *arXiv:1011.2526*.
- [7] I. Benjamini and O. Schramm. Recurrence of distributional limits of finite planar graphs. *Electron. J. Probab.*, 6:no. 23, 13 pp. (electronic), 2001.
- [8] N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1989.
- [9] J. Bouttier, P. Di Francesco, and E. Guitter. Planar maps as labeled mobiles. *Electron. J. Combin.*, 11(1):Research Paper 69, 27 pp. (electronic), 2004.
- [10] D. Burago, Y. Burago, and S. Ivanov. *A course in metric geometry*, volume 33 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001.
- [11] P. Chassaing and B. Durhuus. Local limit of labeled trees and expected volume growth in a random quadrangulation. *Ann. Probab.*, 34(3):879–917, 2006.
- [12] P. Chassaing and G. Schaeffer. Random planar lattices and integrated superBrownian excursion. *Probab. Theory Related Fields*, 128(2):161–212, 2004.
- [13] R. Cori and B. Vauquelin. Planar maps are well labeled trees. *Canad. J. Math.*, 33(5):1023–1042, 1981.
- [14] N. Curien, J.-F. Le Gall, and G. Miermont. The Brownian cactus I. scaling limits of discrete cactuses. *Ann. Inst. H. Poincaré Probab. Statist.* (to appear).

- [15] F. M. Dekking. On transience and recurrence of generalized random walks. *Z. Wahrsch. Verw. Gebiete*, 61(4):459–465, 1982.
- [16] T. Duquesne and J.-F. Le Gall. Random trees, Lévy processes and spatial branching processes. *Astérisque*, (281):vi+147, 2002.
- [17] J. T. Gill and S. Rohde. On the riemann surface type of random planar maps. *arXiv:1101.1320*.
- [18] M. Gromov. Hyperbolic manifolds, groups and actions. In *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978)*, volume 97 of *Ann. of Math. Stud.*, pages 183–213. Princeton Univ. Press, Princeton, N.J., 1981.
- [19] H. Kesten. Subdiffusive behavior of random walk on a random cluster. *Ann. Inst. H. Poincaré Probab. Statist.*, 22(4):425–487, 1986.
- [20] M. Krikun. Local structure of random quadrangulations. *arXiv:0512304*.
- [21] M. Krikun. On one property of distances in the infinite random quadrangulation. *arXiv:0805.1907*.
- [22] M. Krikun. A uniformly distributed infinite planar triangulation and a related branching process. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 307(Teor. Predst. Din. Sist. Komb. i Algoritm. Metody. 10):141–174, 282–283, 2004.
- [23] S. K. Lando and A. Zvonkin. *Graphs on surfaces and their applications*. Springer-Verlag, 2004.
- [24] J.-F. Le Gall. *Spatial branching processes, random snakes and partial differential equations*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 1999.
- [25] J.-F. Le Gall. Random real trees. *Ann. Fac. Sci. Toulouse Math. (6)*, 15(1):35–62, 2006.
- [26] J.-F. Le Gall. Geodesics in large planar maps and in the Brownian map. *Acta Math.*, 205:287–360, 2010.
- [27] J.-F. Le Gall and L. Ménard. Scaling limits for the uniform infinite planar quadrangulation. *arXiv:1005.1738*.
- [28] J.-F. Le Gall and M. Weill. Conditioned Brownian trees. *Ann. Inst. H. Poincaré Probab. Statist.*, 42:455–489, 2006.
- [29] R. Lyons and Y. Peres. *Probability on Trees and Networks*. Current version available at <http://mypage.iu.edu/~rdlyons/>, In preparation.
- [30] L. Ménard. The two uniform infinite quadrangulations of the plane have the same law. *Ann. Inst. H. Poincaré Probab. Statist.*, 46(1):190–208, 2010.
- [31] G. Miermont. Invariance principles for spatial multitype Galton-Watson trees. *Ann. Inst. Henri Poincaré Probab. Stat.*, 44(6):1128–1161, 2008.
- [32] J. Neveu. Arbres et processus de Galton-Watson. *Ann. Inst. H. Poincaré Probab. Statist.*, 22(2):199–207, 1986.
- [33] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*, volume 293 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, third edition, 1999.
- [34] I. Richards. On the classification of noncompact surfaces. *Trans. Amer. Math. Soc.*, 106:259–269, 1963.
- [35] G. Schaeffer. Conjugaison d’arbres et cartes combinatoires aléatoires. phd thesis. 1998.
- [36] W. T. Tutte. A census of planar maps. *Canad. J. Math.*, 15:249–271, 1963.
- [37] A. Zvonkin. Matrix integrals and map enumeration: an accessible introduction. *Math. Comput. Modelling*, 26(8-10):281–304, 1997. Combinatorics and physics (Marseilles, 1995).